



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 31/535, A01N 43/02, 43/40, 43/42, C07D 211/70, 211/72, 217/12, 217/16, 217/18, 217/38, 217/60, 307/02	A1	(11) International Publication Number: WO 99/51241 (43) International Publication Date: 14 October 1999 (14.10.99)
(21) International Application Number: PCT/US99/07760 (22) International Filing Date: 8 April 1999 (08.04.99) (30) Priority Data: 60/081,087 8 April 1998 (08.04.98) US (71) Applicants (for all designated States except US): SMITHK- LINE BEECHAM CORPORATION [US/US]; One Franklin Plaza, Philadelphia, PA 19103 (US). NPS PHARMACEU- TICALS, INC. [US/US]; 420 Chipeta Way, Salt Lake City, UT 84108 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): BHATNAGAR, Pradip, Kumar [US/US]; 300 South Balderston Drive, Exton, PA 19341 (US). CALLAHAN, James, Francis [US/US]; 8214 Jeanes Street, Philadelphia, PA 19111 (US). DEL MAR, Eric, G. [US/US]; 2967 E. St. Mary's Circle, Salt Lake City, UT 84108 (US). LAGO, Maria, Amparo [ES/US]; 701 Pondview Drive, Audubon, PA 19403 (US).		(74) Agents: SIMON, Soma, G. et al.; SmithKline Beecham Corporation, Corporate Intellectual Property, UW2220, 709 Swedeland Road, P.O. Box 1539, King of Prussia, PA 19406-0939 (US). (81) Designated States: AE, AL, AU, BA, BB, BG, BR, CA, CN, CZ, EE, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KP, KR, LC, LK, LR, LT, LV, MG, MK, MN, MX, NO, NZ, PL, RO, SG, SI, SK, SL, TR, TT, UA, US, UZ, VN, YU, ZA, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: CALCILYTIC COMPOUNDS AND METHOD OF USE (57) Abstract Calcilytic compounds and compositions and their use in treating abnormal bone or mineral homeostasis.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

CALCILYTIC COMPOUNDS AND METHOD OF USE

FIELD OF INVENTION

The present invention relates to novel calcilytic compounds,
5 pharmaceutical compositions containing these compounds and their use as calcium
receptor antagonists.

In mammals, extracellular Ca^{2+} is under rigid homeostatic control and
regulates various processes such as blood clotting, nerve and muscle excitability,
and proper bone formation. Extracellular Ca^{2+} inhibits the secretion of parathyroid
10 hormone ("PTH") from parathyroid cells, inhibits bone resorption by osteoclasts,
and stimulates secretion of calcitonin from C-cells. Calcium receptor proteins
enable certain specialized cells to respond to changes in extracellular Ca^{2+}
concentration.

PTH is the principal endocrine factor regulating Ca^{2+} homeostasis in the
15 blood and extracellular fluids. PTH, by acting on bone and kidney cells, increases
the level of Ca^{2+} in the blood. This increase in extracellular Ca^{2+} then acts as a
negative feedback signal, depressing PTH secretion. The reciprocal relationship
between extracellular Ca^{2+} and PTH secretion forms an important mechanism
maintaining bodily Ca^{2+} homeostasis.

20 Extracellular Ca^{2+} acts directly on parathyroid cells to regulate PTH
secretion. The existence of a parathyroid cell surface protein which detects
changes in extracellular Ca^{2+} has been confirmed. See Brown *et al.*, *Nature*
366:574, 1993. In parathyroid cells, this protein, the calcium receptor, acts as a
receptor for extracellular Ca^{2+} , detects changes in the ion concentration of
25 extracellular Ca^{2+} , and initiates a functional cellular response, PTH secretion.

Extracellular Ca^{2+} influences various cell functions, reviewed in Nemeth
et al., *Cell Calcium* 11:319, 1990. For example, extracellular Ca^{2+} plays a role in
parafollicular (C-cells) and parathyroid cells. See Nemeth, *Cell Calcium* 11:323,
1990. The role of extracellular Ca^{2+} on bone osteoclasts has also been studied. See
30 Zaidi, *Bioscience Reports* 10:493, 1990.

Various compounds are known to mimic the effects of extra-cellular Ca^{2+}
on a calcium receptor molecule. Calcilytics are compounds able to inhibit calcium
receptor activity, thereby causing a decrease in one or more calcium receptor

activities evoked by extracellular Ca^{2+} . Calcilytics are useful as lead molecules in the discovery, development, design, modification and/or construction of useful calcium modulators which are active at Ca^{2+} receptors. Such calcilytics are useful in the treatment of various disease states characterized by abnormal levels of one or more components, e.g., polypeptides such as hormones, enzymes or growth factors, the expression and/or secretion of which is regulated or affected by activity at one or more Ca^{2+} receptors. Target diseases or disorders for calcilytic compounds include diseases involving abnormal bone and mineral homeostasis.

Abnormal calcium homeostasis is characterized by one or more of the following activities: an abnormal increase or decrease in serum calcium; an abnormal increase or decrease in urinary excretion of calcium; an abnormal increase or decrease in bone calcium levels (for example, as assessed by bone mineral density measurements); an abnormal absorption of dietary calcium; an abnormal increase or decrease in the production and/or release of messengers which affect serum calcium levels such as PTH and calcitonin; and an abnormal change in the response elicited by messengers which affect serum calcium levels.

Thus, calcium receptor antagonists offer a unique approach towards the pharmacotherapy of diseases associated with abnormal bone or mineral homeostasis, such as hypoparathyroidism, osteosarcoma, periodontal disease, fracture healing, osteoarthritis, rheumatoid arthritis, Paget's disease, humoral hypercalcemia associated with malignancy and fracture healing, and osteoporosis.

SUMMARY OF THE INVENTION

The present invention comprises novel calcium receptor antagonists represented by Formula (I) hereinbelow and their use in the treatment of a variety of diseases associated with abnormal bone or mineral homeostasis, including but not limited to hypoparathyroidism, osteosarcoma, periodontal disease, fracture healing, osteoarthritis, rheumatoid arthritis, Paget's disease, humoral hypercalcemia associated with malignancy and fracture healing, and osteoporosis.


The present invention further provides a method for antagonizing calcium receptors in an animal, including humans, which comprises administering to an

animal in need thereof an effective amount of a compound of Formula (I), indicated hereinbelow.

The present invention further provides a method for increasing serum parathyroid levels in an animal, including humans, which comprises administering to an animal in need thereof an effective amount of a compound of Formula (I), indicated hereinbelow.

DETAILED DESCRIPTION OF THE INVENTION

The compounds of the present invention are selected from Formula (I) hereinbelow:



Formula (I)

wherein:

Y₁ is a covalent bond, alkylene or alkenylene of up to 4 carbon atoms, unsubstituted or substituted by C₁₋₄ alkyl or O;

Y₂ is methylene, unsubstituted or substituted by C₁₋₄ alkyl or haloalkyl;

Y₃ is covalent bond or O, S, N-R^{IV} or C₁₋₄ alkylene-O, C₁₋₄ alkylene-S, C₁₋₄ alkylene-N-R^{IV};

R^{IV} is selected from the group consisting of H, C₁₋₄ alkyl, C₃₋₆ cycloalkyl;

R₃ and R₄ are, independently, methyl or ethyl, or, together, form cyclopropyl;

R₅ is heteroaryl or fused heteroaryl; wherein the hetero-ring contains N, O or S, and is aromatic, dihydro or tetrahydro, unsubstituted or substituted with any

substituents being selected from the group consisting of OH, OCH₃, CH(CH₃)₂, halogen, C₁₋₄ alkyl, C₁₋₄ alkoxy, C₃₋₆ cycloalkyl, OSO₂R^{IV}, CN, NO₂, OCF₃, CF₃, CH₂CF₃, (CH₂)_n CO₂H, (CH₂)_n CO₂R^{IV}, and O-(CH₂)_n CO₂R^{IV};

n is an integer from 0 to 3;

G is a covalent bond, CHR₆ or C-R₆, wherein R₆ is H, OH or O (forming a ketone);

R₇ is H, OH, or O-C₁₋₄ alkyl;


R₈ is H or C₁₋₄ alkyl; or R₇ and R₈ together form a ketone;

A and B are, independently, selected from the group consisting of a bond, CH₂,


NH, O, S and C=O, provided that either A or B is selected from CH₂ and NH; or A

5 and B together form a bond; or the A-B moiety is represented by CH=CH or C≡C;


X is selected from sub formulas (Ia) to (Ie) hereinbelow:




(Ia)




(Ib)



(Ic)



(Id)



(Ie)

5

wherein

- W is selected from the group consisting of R_1 , SO_2R_1 , $C(O)R_1$, $SO_2NR_1R_1'$, $C(O)NR_1R_1'$, $C(O)OR_1$, SO_3R_1' , wherein R_1 and R_1' are independently selected from the group consisting of hydrogen, C_{1-4} alkyl, C_{3-6} cycloalkyl, C_{2-5} alkenyl, C_{2-5} alkynyl, heterocycloalkyl, aryl and aryl C_{1-4} alkyl; or R_1 and R_1' together form a 3 to 7 membered optionally substituted heterocyclic ring; wherein any substituents are selected from the group consisting of CN, aryl, CO_2R , CO_2NHR , OH, OR, NH_2 , halo, CF_3 , OCF_3 and NO_2 ; wherein R represents C_{1-4} alkyl, or C_{3-6} cycloalkyl;
- 15 X_1 is selected from the group consisting of CN, NO_2 , Cl, F, Br, I, H, R' , OR', CF_3 , OCF_3 and OSO_2R' , wherein R' represents C_{1-4} alkyl, or C_{3-6} cycloalkyl; X_2 , X_3 and X_4 are, independently, selected from the group consisting of CN, NO_2 , Cl, F, Br, I, H, R'' , OR'', CF_3 , OCF_3 and OSO_2R'' , provided that either X_1 or X_3 is H, wherein R'' is C_{1-4} alkyl or haloalkyl; or X_1 and X_2 together form an aryl or heteroaryl ring, substituted or unsubstituted; wherein the heteroatom is selected from N, S and O; and any substituents are selected from the group consisting of halo, C_{1-4} alkyl, OCF_3 , CF_3 , OMe, CN, OSO_2R' and NO_2 ; or X_3 and X_4 independently represent $C(O)R_1$; and
- 20


- R_2 is selected from the group consisting of hydrogen, C_{1-4} alkyl, C_{3-6} cycloalkyl, C_{2-5} alkenyl, C_{2-5} alkynyl, heterocycloalkyl aryl and aryl- C_{1-4} alkyl;
- X_1 is selected from the group consisting of CN, NO_2 , Cl, F, Br, I, H, R, OR, CF_3 , OCF_3 and OSO_2R , wherein R represents C_{1-4} alkyl, or C_{3-6} cycloalkyl;
- 5 X_2 , X_3 and X_4 are, independently, selected from the group consisting of CN, NO_2 , Cl, F, Br, I, H, R', OR', CF_3 , OCF_3 and OSO_2R' , provided that either X_1 or X_3 is H, wherein R' is C_{1-4} alkyl or haloalkyl; or X_1 and X_2 together form an aryl or heteroaryl ring, substituted or unsubstituted; wherein the heteroatom is selected from N, S and O and any substituents are selected from the group
- 10 consisting of halo, C_{1-4} alkyl, OCF_3 , CF_3 , OMe, CN, OSO_2-C_{1-4} alkyl, OSO_2-C_{3-6} cycloalkyl and NO_2 ;
- or X_3 and X_4 independently represent $C(O)R_1$; and
- R_1 and R_2 are, independently, selected from the group consisting of hydrogen, C_{1-4} alkyl, C_{3-6} cycloalkyl, C_{2-5} alkenyl, C_{2-5} alkynyl, heterocycloalkyl and
- 15 aryl; or R_1 and R_2 together form a 3 to 7 membered optionally substituted heterocyclic ring; wherein any substituents are selected from the group consisting of CN, aryl, CO_2R , CO_2NHR , OH, OR, NH_2 , halo, CF_3 , OCF_3 and NO_2 ;
- wherein R represents C_{1-4} alkyl, or C_{3-6} cycloalkyl;
- X_1 is selected from the group consisting of CN, NO_2 , Cl, F, Br, I, H, R, OR, CF_3 ,
- 20 OCF_3 and OSO_2R , wherein R represents C_{1-4} alkyl, or C_{3-6} cycloalkyl;
- X_2 , X_3 , and X_4 are, independently, selected from the group consisting of CN, NO_2 , Cl, F, Br, I, H, R', OR', CF_3 , OCF_3 and OSO_2R' , provided that either X_1 or X_3 is H, wherein R' is C_{1-4} alkyl or haloalkyl;
- or X_1 and X_2 together form an aryl or heteroaryl ring, substituted or
- 25 unsubstituted; wherein the heteroatom is selected from N, S and O and the substituents are selected from the group consisting of halo, C_{1-4} alkyl, OCF_3 , CF_3 , OMe, CN, OSO_2-C_{1-4} alkyl, OSO_2-C_{3-6} cycloalkyl and NO_2 ;
- or X_3 and X_4 independently represent $C(O)R_1$;
- R_1 and R_2 are, independently, selected from the group consisting of hydrogen,
- 30 C_{1-4} alkyl, C_{3-6} cycloalkyl, C_{2-5} alkenyl, C_{2-5} alkynyl, heterocycloalkyl and aryl;
- or R_1 and R_2 together form a 3 to 7 membered optionally substituted heterocyclic ring; wherein the substituents are selected from the group consisting of

- CN, aryl, CO₂R", CO₂NHR", OH, OR", NH₂, halo, CF₃, OCF₃ and NO₂; wherein R" represents C₁₋₄ alkyl, or C₃₋₆ cycloalkyl;
- D is selected from the group consisting of H, CN, NO₂, Cl, F, Br, I, R, OR, SR, CF₃, OCF₃ and OSO₂R, wherein R represents C₁₋₄ alkyl, C₃₋₆ cycloalkyl, or C₁₋
- 5 10 aryl or heteroaryl wherein the heteroatom is selected from N, S and O and substituents are selected from the group consisting of halo, C₁₋₄ alkyl, OCF₃, CF₃, OMe, CN, OSO₂-C₁₋₄ alkyl, OSO₂-C₃₋₆ cycloalkyl and NO₂;
- n is the integer 1 or 2;
- 10 each E is independently C or N, provided that no more than two E moieties are N; further provided that when n is 2, each E is C;
- a and b are optionally present bonds;
- R₁^{iv} is selected from the group consisting of (CH₂)_nCO₂R', (CH₂)_nCO₂H, (CH₂)_nCONR'₂, (CH₂)_nCH₂OR', OR', SR', CN, NO₂, Cl, F, Br, I, H, CF₃, OCF₃,
- 15 OSO₂R', R' and H; wherein R' represents C₁₋₄ alkyl, or C₃₋₆ cycloalkyl; or R₁^{iv} is O, forming a ketone such that Y R₁^{iv} represents -C=O;
- R₂^{iv} is selected from the group consisting of hydrogen, CN, NO₂, Cl, F, Br, I, H, R", OR", CF₃, OCF₃, and OSO₂R", wherein R" represents C₁₋₄ alkyl, or C₃₋₆ cycloalkyl.
- 20 Y is selected from the group consisting of C, CH, O, N and S; provided that when Y is S, R₁^{iv} is O or not present; further provided that when Y is O, R₁^{iv} is not present;
- X' is selected from the group consisting of CH₂, NH, O and S.
- R₉ is selected from the group consisting of O-alkyl, O-CH₂-aryl, and O- aryl;
- 25 X₁ⁱⁱⁱ is selected from the group consisting of CN, NO₂, Cl, F, Br, I, H, R, OR, CF₃, OCF₃ and OSO₂R, wherein R represents C₁₋₄ alkyl, or C₃₋₆ cycloalkyl;
- X₂ⁱⁱⁱ, X₃ⁱⁱⁱ, and X₄ⁱⁱⁱ are, independently, selected from the group consisting of CN, NO₂, Cl, F, Br, I, H, R', OR', CF₃, OCF₃ and OSO₂R', provided that either Xⁱⁱⁱ₁ or Xⁱⁱⁱ₃ is H, wherein R' is C₁₋₄ alkyl or haloalkyl;
- 30 or X₁ⁱⁱⁱ and X₂ⁱⁱⁱ together form an aryl or heteroaryl ring, substituted or unsubstituted; wherein the heteroatom is selected from N, S and O and the substituents are selected from the group consisting of halo, C₁₋₄ alkyl, OCF₃, CF₃, OMe, CN, OSO₂-C₁₋₄ alkyl, OSO₂-C₃₋₆ cycloalkyl and NO₂;

or X_2^m and X_4^m independently represent $C(O)R_1$;

and pharmaceutically acceptable salts and complexes thereof.

Preferably, the compounds of the present invention have a structure according to Formula (II):



Formula (II)

wherein:

- R_5 is heteroaryl or fused heteroaryl; wherein the hetero-ring contains N, O or S, and is aromatic, dihydro or tetrahydro, unsubstituted or substituted with any substituents being selected from the group consisting of OH, OCH_3 , $CH(CH_3)_2$, halogen, C_{1-4} alkyl, C_{1-4} alkoxy, C_{3-6} cycloalkyl, OSO_2R^{IV} , CN, NO_2 , OCF_3 , CF_3 , CH_2CF_3 , $(CH_2)_n CO_2H$, $(CH_2)_n CO_2R^{IV}$, and $O-(CH_2)_n CO_2R^{IV}$; and A and B are, independently, selected from the group consisting of a bond, CH_2 , NH, O, S and C=O, provided that either A or B is selected from CH_2 and NH; or A and B together form a bond; or the A-B moiety is represented by $CH=CH$ or $C\equiv C$.

- More preferably, R_5 is heteroaryl or fused heteroaryl, wherein the hetero-ring contains N, O or S and is aromatic, dihydro or tetrahydro, unsubstituted or substituted with any substituents being selected from the group consisting of OCH_3 , halogen, C_{1-4} alkyl, , CN, NO_2 , OCF_3 , CF_3 , CH_2CF_3 ;

R_6 is H; and

A and B are, independently, selected from the group consisting of a bond, CH_2 , NH, O, S and C=O, provided that either A or B is selected from CH_2 and NH, or A and B together form a bond.

- Most preferably, R_5 is heteroaryl or fused heteroaryl, wherein the hetero-ring contains N, O or S and is aromatic, dihydro or tetrahydro, unsubstituted or substituted with any substituents being selected from the group consisting of OCH_3 , halogen, C_{1-4} alkyl, , CN, NO_2 , OCF_3 , CF_3 , CH_2CF_3 ;

R_6 is H; and

- A and B are, independently, selected from the group consisting of a bond, CH_2 , O, or A and B together form a bond.

In sub-formula (Ia), preferably, X_1 is selected from the group consisting of CN, NO_2 , Cl, F, Br, I and H. Preferably, X_2 , X_3 and X_4 are, independently, selected from the group consisting of Cl, F, Br, I and H, provided X_1 and X_3 is H. Preferably, R_1 , R_1' and R_2 are, independently, selected from the group consisting of

5 C_{1-4} alkyl, C_{3-6} cycloalkyl, heterocycloalkyl, aryl or arylalkyl.

In sub-formula (Ia), more preferably, R_1 , R_1' and R_2 are, independently, H, alkyl, or aryl. More preferably, X_1 is selected from the group consisting of CN, NO_2 , Cl, F, Br, I and H. More preferably, X_2 , X_3 and X_4 are, independently, selected from the group consisting of Cl, F, Br, I and H provided X_1 and X_3 is H.

10 In sub-formula (Ia), more preferably still, R_1 , R_1' and R_2 are, independently, C_{1-4} alkyl, or aryl. More preferably still, X_1 is CN, NO_2 , or Cl. More preferably still, X_2 is Cl, F or H. More preferably still, X_3 and X_4 are H.

In sub-formula (Ia), most preferably, X_1 is CN, or NO_2 . Most preferably, X_2 is Cl.

15 In sub-formula (Ib), preferably, X_1'' is selected from the group consisting of CN, NO_2 , Cl, F, Br, I and H. Preferably, X_2'' , X_3'' and X_4'' are, independently, selected from the group consisting of Cl, F, Br, I and H. Preferably, R_1'' and R_2'' are, independently, selected from the group consisting of C_{1-4} alkyl, C_{3-6} cycloalkyl, heterocycloalkyl or aryl; or R_1'' and R_2'' together form an optionally

20 substituted 3-7 membered ring, optionally containing an additional heteroatom selected from O, S, and N.

In sub-formula (Ib), more preferably, R_1'' and R_2'' are, independently, H, C_{1-4} alkyl, or aryl; or R_1'' and R_2'' together form an optionally substituted 4-7 membered ring, optionally containing a heteroatom selected from O, S, and N.

25 More preferably, X_1'' is selected from the group consisting of CN, NO_2 , Cl, F, Br, I and H. More preferably, X_2'' is selected from the group consisting of Cl, F, Br, I and H.

In sub-formula (Ib), more preferably still, R_1'' and R_2'' are, independently, C_{1-4} alkyl, or aryl; or R_1'' and R_2'' together form a 4-7 membered ring as described

30 hereinabove. More preferably still, X_1'' is CN, NO_2 , or Cl. More preferably still, X_2'' is Cl, F or H.

In sub-formula (Ib), most preferably, R_1'' and R_2'' together form a 4-7 membered ring as described hereinabove. Most preferably, X_1'' is CN, or NO_2 . Most preferably, X_2'' is Cl.

In sub-formula (Ic), preferably, X_1''' is selected from the group consisting of CN, NO_2 , Cl, F, Br, I and H. Preferably, X_2''' , X_3''' and X_4''' are, independently, selected from the group consisting of Cl, F, Br, I and H. provided either X_1''' or X_3''' is H. Preferably, R_1''' and R_2''' are, independently, selected from the group consisting of C_{1-4} alkyl, C_{3-6} cycloalkyl, heterocycloalkyl or aryl; or R_1''' and R_2''' together form an optionally substituted 3-7 membered ring, optionally containing an additional heteroatom selected from O, S, and N.

In sub-formula (Ic), more preferably, R_1''' and R_2''' are, independently, H, C_{1-4} alkyl, or aryl; or R_1''' and R_2''' together form an optionally substituted 4-7 membered ring, optionally containing a heteroatom selected from O, S, and N. More preferably, X_1''' is selected from the group consisting of CN, NO_2 , Cl, F, Br, I and H. More preferably, X_2''' , X_3''' and X_4''' are, independently, selected from the group consisting of Cl, F, Br, I and H provided either X_1''' or X_3''' is H.

In sub-formula (Ic), more preferably still, R_1''' and R_2''' are, independently, C_{1-4} alkyl, or aryl; or R_1''' and R_2''' together form a 4-7 membered ring as described hereinabove. More preferably still, X_1''' is CN, NO_2 , or Cl. More preferably still, X_2''' is Cl or H. More preferably still, X_3''' and X_4''' are H.

In sub-formula (Ic), most preferably, R_1''' and R_2''' together form a 4-7 membered ring as described hereinabove. Most preferably, X_1''' is CN or NO_2 . Most preferably, X_2''' is Cl.

In sub-formula (Id), preferably, each D is selected from the group consisting of F, Br, Cl, I, R, OR, SR, and H. Preferably, R_1^{IV} is selected from the group consisting of $(CH_2)_nCO_2R'$, $(CH_2)_nCO_2H$, $(CH_2)_nCONR'_2$, $(CH_2)_nCH_2OR'$, OR', SR', R' and H; wherein R' is as R hereinabove; or R_1^{IV} is O, forming a ketone such that $Y R_1^{IV}$ represents $-C=O$. Preferably, R_2^{IV} is selected from the group consisting of hydrogen, CN, NO_2 , Cl, Br, F and I;

In sub-formula (Id), more preferably, n is 0. More preferably, each E is C. More preferably, X' is CH_2 , O, or NH. More preferably, Y is C or N. More preferably, R_1^{IV} is CH_2CO_2R' , SR', or O forming a ketone.

In subformula (Id), more preferably still, X' is CH₂ or O. More preferably still, R₁^{iv} is CH₂CO₂R' or SR'. More preferably still, R₂^{iv} is H, CN, or NO₂.

In subformula (Id), most preferably, X' is CH₂. Most preferably, Y is C. Most preferably, R₂^{iv} is CN or NO₂.

- 5 In subformula (Ie), preferably R₉ is selected from the group consisting of O-(CH₂)_n-aryl, and O- aryl;
 X₁ⁱⁱⁱ is selected from the group consisting of CN, NO₂, Cl, F, Br, H, R, and OSO₂R, wherein R represents C₁₋₄ alkyl, or C₃₋₆ cycloalkyl;
 X₂ⁱⁱⁱ, X₃ⁱⁱⁱ, and X₄ⁱⁱⁱ are, independently, selected from the group consisting of CN,
 10 NO₂, Cl, F, Br, H, and OSO₂R', provided that either Xⁱⁱⁱ₁ or Xⁱⁱⁱ₃ is H, wherein R' is C₁₋₄ alkyl or haloalkyl;
 or X₁ⁱⁱⁱ and X₂ⁱⁱⁱ together form an aryl or heteroaryl ring, substituted or unsubstituted; wherein the heteroatom is selected from N, S and O and the substituents are selected from the group consisting of halo, C₁₋₄ alkyl, OCF₃, CF₃,
 15 OMe, CN, OSO₂-C₁₋₄ alkyl, OSO₂-C₃₋₆ cycloalkyl and NO₂;
 or X₂ⁱⁱⁱ and X₄ⁱⁱⁱ independently represent C(O)R₁;

- In subformula (Ie), more preferably R₉ is selected from the group consisting of O-(CH₂)_n-aryl, and O- aryl;
 X₁ⁱⁱⁱ is selected from the group consisting of CN, NO₂, and Cl
 20 X₂ⁱⁱⁱ, X₃ⁱⁱⁱ, and X₄ⁱⁱⁱ are, independently, selected from the group consisting of Cl, F, and H, provided that either Xⁱⁱⁱ₁ or Xⁱⁱⁱ₃ is H,
 or X₂ⁱⁱⁱ and X₄ⁱⁱⁱ independently represent C(O)R₁;

- In subformula (Ie), most preferably R₉ is selected from the group consisting of O-(CH₂)_n-aryl, and O- aryl;
 25 X₁ⁱⁱⁱ is CN or NO₂,
 X₂ⁱⁱⁱ is Cl, X₃ⁱⁱⁱ and X₄ⁱⁱⁱ are, independently F, and H.

Preferred heteroaryls useful in the present invention include unsubstituted and substituted quinolines, isoquinolines, benzofurans, dihydrobenzofurans, benzothiophenes, dihydrobenzothiophenes and pyridines.

- 30 As used herein "cycloalkyl" refers to optionally substituted 3-7 membered carbocyclic rings wherein any substituents are selected from the group consisting of, F, Cl, Br, I, N(R₁)₂, SR₁ and OR₁, unless otherwise indicated.

As used herein "heterocycloalkyl" refers to optionally substituted 4, 5, 6 or 7 membered heterocyclic rings containing 1 to 2 heteroatoms selected from N, O, and S.

As used herein, "aryl" refers to an optionally substituted aromatic group with at least one ring having a conjugated pi-electron system, containing up to two conjugated or fused ring systems. Aryl includes carbocyclic aryl, and biaryl groups, all of which may be optionally substituted. Preferred aryl include phenyl and naphthyl. More preferred aryl include phenyl. Preferred substituents are selected from the group consisting of halo, C₁₋₄ alkyl, OCF₃, CF₃, OMe, CN, OSO₂ R and NO₂, wherein R represents C₁₋₄ alkyl or C₃₋₆ cycloalkyl.

As used herein, "acyl" refers to C₁₋₄ alkylcarbonyl.

As used herein, "alkenyl" refers to an optionally substituted hydrocarbon group containing at least one carbon-carbon double bond and containing upto 5 carbon atoms joined together. The alkenyl hydrocarbon chain may be straight, branched or cyclic. Any substituents are selected from the group consisting of halo, C₁₋₄ alkyl, OCF₃, CF₃, OMe, CN, OSO₂ R and NO₂, wherein R represents C₁₋₄ alkyl or C₃₋₆ cycloalkyl.

As used herein, "alkynyl" refers to an optionally substituted hydrocarbon group containing at least one carbon-carbon triple bond between the carbon atoms and containing up to 5 carbon atoms joined together. The alkynyl hydrocarbon group may be straight-chained, branched or cyclic. Any substituents are selected from the group consisting of halo, C₁₋₄ alkyl, OCF₃, CF₃, OMe, CN, OSO₂ R and NO₂, wherein R represents C₁₋₄ alkyl or C₃₋₆ cycloalkyl.

The compounds of the present invention may contain one or more asymmetric carbon atoms and may exist in racemic and optically active forms. All of these compounds and diastereomers are contemplated to be within the scope of the present invention.

Preferred compounds of the present invention are selected from the group consisting of:

- (R)-N-[2-Hydroxy-3-(3-chloro-2-cyanophenoxy)propyl]-1,1-dimethyl-2-(2,3-dihydrobenzo[b]furan-5yl)ethylamine;
(R)-N-[2-Hydroxy-3-(3-chloro-2-cyanophenoxy)propyl]-1,1-dimethyl-2-(quinolin-3-yl)ethylamine;

- (R)-N-[2-Hydroxy-3-(3-chloro-2-cyanophenoxy)propyl]-1,1-dimethyl-2-(quinolin-2-yl)ethylamine;
- (R)-N-[2-Hydroxy-3-(3-chloro-2-cyanophenoxy)propyl]-1,1-dimethyl-2-(isoquinolin-3-yl)ethylamine;
- 5 (R)-N-[2-Hydroxy-3-(3-chloro-2-cyanophenoxy)propyl]-1,1-dimethyl-4-(2-pyridyl)butylamine;
- (R)-N-[2-Hydroxy-3-(3-chloro-2-cyano-4-morpholinosulfonamidophenoxy)propyl]-1,1-dimethyl-4-(2-pyridyl)butylamine;
- (R)-N-[2-Hydroxy-3-(3-chloro-2-cyanophenoxy)propyl]-1,1-dimethyl-4-(3-pyridyl)butylamine;
- 10 (R)-N-[2-Hydroxy-3-(3-chloro-2-cyano-4-morpholinosulfonamidophenoxy)propyl]-1,1-dimethyl-4-(3-pyridyl)butylamine;
- (R)-N-[2-Hydroxy-3-(3-chloro-2-cyanophenoxy)propyl]-1,1-dimethyl-4-(4-carbethoxyphenyl)butylamine;
- 15 (R)-N-[2-Hydroxy-3-(3-chloro-2-cyanophenoxy)propyl]-1,1-dimethyl-2-(4-ethylpyrid-2-yl)ethylamine;
- (R)-N-[2-Hydroxy-3-(3-chloro-2-cyanophenoxy)propyl]-1,1-dimethyl-2-benzamidoethylamine;
- (R)-N-[2-Hydroxy-3-(3-chloro-2-cyanophenoxy)propyl]-4-phenylbutylamine;
- 20 (R)-N-[2-Hydroxy-3-(3-chloro-2-cyanophenoxy)propyl]-1,1-dimethyl-4-phenylbut-2-ynylamine;
- and pharmaceutically acceptable salts and complexes thereof. Preferred salts include hydrochloride and dihydrochloride.

More preferred compounds useful in the present invention include:

- 25 (R)-N-[2-Hydroxy-3-(3-chloro-2-cyanophenoxy)propyl]-1,1-dimethyl-2-(2,3-dihydrobenzo[b]furan-5yl)ethylamine;
- (R)-N-[2-Hydroxy-3-(3-chloro-2-cyanophenoxy)propyl]-1,1-dimethyl-2-(quinolin-3-yl)ethylamine;
- (R)-N-[2-Hydroxy-3-(3-chloro-2-cyanophenoxy)propyl]-1,1-dimethyl-2-(quinolin-2-yl)ethylamine;
- 30 (R)-N-[2-Hydroxy-3-(3-chloro-2-cyanophenoxy)propyl]-1,1-dimethyl-2-(isoquinolin-3-yl)ethylamine;

(R)-N-[2-Hydroxy-3-(3-chloro-2-cyanophenoxy)propyl]-1,1-dimethyl-4-(2-pyridyl)butylamine;

(R)-N-[2-Hydroxy-3-(3-chloro-2-cyano-4-morpholinosulfonamidophenoxy)propyl]-1,1-dimethyl-4-(2-pyridyl)butylamine;

5 (R)-N-[2-Hydroxy-3-(3-chloro-2-cyanophenoxy)propyl]-1,1-dimethyl-4-(3-pyridyl)butylamine;

(R)-N-[2-Hydroxy-3-(3-chloro-2-cyano-4-morpholinosulfonamidophenoxy)propyl]-1,1-dimethyl-4-(3-pyridyl)butylamine;

10 (R)-N-[2-Hydroxy-3-(3-chloro-2-cyanophenoxy)propyl]-1,1-dimethyl-2-(4-ethylpyrid-2-yl)ethylamine;

(R)-N-[2-Hydroxy-3-(3-chloro-2-cyanophenoxy)propyl]-1,1-dimethyl-2-benzamidoethylamine;

(R)-N-[2-Hydroxy-3-(3-chloro-2-cyanophenoxy)propyl]-4-phenylbutylamine; and pharmaceutically acceptable salts and complexes thereof.

15 The most preferred compounds useful in the present invention include:

(R)-N-[2-Hydroxy-3-(3-chloro-2-cyanophenoxy)propyl]-1,1-dimethyl-2-(2,3-dihydrobenzo[b]furan-5yl)ethylamine;

(R)-N-[2-Hydroxy-3-(3-chloro-2-cyanophenoxy)propyl]-1,1-dimethyl-2-(quinolin-3-yl)ethylamine;

20 (R)-N-[2-Hydroxy-3-(3-chloro-2-cyanophenoxy)propyl]-1,1-dimethyl-2-(quinolin-2-yl)ethylamine;

(R)-N-[2-Hydroxy-3-(3-chloro-2-cyanophenoxy)propyl]-1,1-dimethyl-2-(isoquinolin-3-yl)ethylamine;

25 (R)-N-[2-Hydroxy-3-(3-chloro-2-cyanophenoxy)propyl]-1,1-dimethyl-4-(2-pyridyl)butylamine;

(R)-N-[2-Hydroxy-3-(3-chloro-2-cyano-4-morpholinosulfonamidophenoxy)propyl]-1,1-dimethyl-4-(2-pyridyl)butylamine;

(R)-N-[2-Hydroxy-3-(3-chloro-2-cyanophenoxy)propyl]-1,1-dimethyl-4-(3-pyridyl)butylamine;

30 (R)-N-[2-Hydroxy-3-(3-chloro-2-cyano-4-morpholinosulfonamidophenoxy)propyl]-1,1-dimethyl-4-(3-pyridyl)butylamine;

(R)-N-[2-Hydroxy-3-(3-chloro-2-cyanophenoxy)propyl]-1,1-dimethyl-2-(4-ethylpyrid-2-yl)ethylamine;

and pharmaceutically acceptable salts and complexes thereof.

Pharmaceutically acceptable salts are non-toxic salts in the amounts and concentrations at which they are administered.


Pharmaceutically acceptable salts include acid addition salts such as those
 5 containing sulfate, hydrochloride, fumarate, maleate, phosphate, sulfamate, acetate, citrate, lactate, tartrate, methanesulfonate, ethanesulfonate, benzenesulfonate, *p*-toluenesulfonate, cyclohexylsulfamate and quinate. A preferred salt is a hydrochloride. Pharmaceutically acceptable salts can be obtained from acids such
 10 as hydrochloric acid, maleic acid, sulfuric acid, phosphoric acid, sulfamic acid, acetic acid, citric acid, lactic acid, tartaric acid, malonic acid, methanesulfonic acid, ethanesulfonic acid, benzenesulfonic acid, *p*-toluenesulfonic acid, cyclohexylsulfamic acid, fumaric acid, and quinic acid.

Pharmaceutically acceptable salts also include basic addition salts such as those containing benzathine, chlorprocaine, choline, diethanolamine,
 15 ethylenediamine, meglumine, procaine, aluminum, calcium, lithium, magnesium, potassium, sodium, ammonium, alkylamine, and zinc, when acidic functional groups, such as carboxylic acid or phenol are present.

The present invention provides compounds of Formula (I) above, which can be prepared using standard techniques. An overall strategy for preparing
 20 preferred compounds described herein can be carried out as described in this section. The examples which follow illustrate the synthesis of specific compounds. Using the protocols described herein as a model, one of ordinary skill in the art can readily produce other compounds of the present invention.


25

Scheme 1




30

Scheme 2




5

Scheme 3



10

Scheme 4



5

General Preparation

A general procedure used to synthesize many of the compounds can be carried out as described in Scheme 1: A solution of aryl alcohol in acetone was treated with an appropriate base such as K_2CO_3 , heated for 15 min. R-glycidyl nosylate was added and the reaction continued overnight to give the corresponding glycidyl ether (Scheme 1). In the case of an alkyl alcohol, a stronger base, e.g. NaH in DMF was used. This method can also be used for aryl alcohols. A solution of the substituted glycidyl ether and excess amine (typically 1,1-dimethyl-2-(4-methyloxyphenyl)ethylamine) in absolute ethanol, acetonitrile, THF or any other similar solvent in the presence of a suitable catalyst such as LiClO_4 is stirred overnight at reflux. The product is purified by normal phase chromatography. Hydrochloride salts are prepared by treatment of the corresponding free base with HCl either in gas phase or 4M dioxane solution, or any other standard method.

The synthesis of various corresponding amines is described in Scheme 2, 3, 4 and 5. The synthesis of 3-(2-amino-2-methylpropyl)quinoline illustrates the general procedure to obtain these amines, and it is described in Scheme 2. The reduction of the oxime obtained from 3-quinolinecarboxaldehyde leads to the corresponding benzylic amine. Reaction of the aforementioned amine with 2,4,6-triphenylpyrylium tetrafluoroborate followed by nucleophilic displacement of the

pyridinium salt thus formed with the anion of 2-nitropropane, leads to the formation of the corresponding nitro compound which, after reduction, leads to the title compound.

The synthesis of 2-(4-amino-4-methylpentyl)pyridine illustrates the general procedure to obtain pentyl amines, and it is described in Scheme 3. The Curtius rearrangement of 2,2-dimethyl-4-pentenoic acid leads to the corresponding Cbz protected amine. Addition of 9-BBN to the terminal olefin of the protected amine leads to the corresponding boronate. Palladium catalyzed coupling reaction between the boronate and the corresponding aryl bromide (2-bromopyridine in Scheme 3) leads to the formation of the corresponding amine after the removal of the protecting group.

The synthesis of 5-(2-amino-2-methylpropyl)-2,3-dihydrobenzo[b]furan illustrates the general procedure to obtain these amines, and it is described in Scheme 4. Wittig reaction between 2,3-dihydrobenzo[b]furan-5-carboxaldehyde and the anion formed from isopropyltriphenylphosphonium leads to the corresponding olefin. Ritter reaction on the olefin followed by hydrolysis leads to the corresponding amine.

Nuclear magnetic resonance spectra were recorded at either 250 or 400 MHz using, respectively, a Bruker AM 250 or Bruker AC 400 spectrometer. CDCl₃ is deuteriochloroform, DMSO-d₆ is hexadeuteriodimethylsulfoxide, and CD₃OD is tetradeuteriomethanol. Chemical shifts are reported in parts per million (•) downfield from the internal standard tetramethylsilane. Abbreviations for NMR data are as follows: s=singlet, d=doublet, t=triplet, q=quartet, m=multiplet, dd=doublet of doublets, dt=doublet of triplets, app=apparent, br=broad. J indicates the NMR coupling constant measured in Hertz. Continuous wave infrared (IR) spectra were recorded on a Perkin-Elmer 683 infrared spectrometer, and Fourier transform infrared (FTIR) spectra were recorded on a Nicolet Impact 400 D infrared spectrometer. IR and FTIR spectra were recorded in transmission mode, and band positions are reported in inverse wavenumbers (cm⁻¹). Mass spectra were taken on either VG 70 FE, PE Syx API III, or VG ZAB HF instruments, using fast atom bombardment (FAB) or electrospray (ES) ionization techniques. LC/MS/MS was obtained on a Perkin Elmer Sciex API 365 Instrument. Elemental analyses were obtained using a Perkin-Elmer 240C elemental analyzer. Melting

points were taken on a Thomas-Hoover melting point apparatus and are uncorrected. All temperatures are reported in degrees Celsius.

Analtech Silica Gel GF and E. Merck Silica Gel 60 F-254 thin layer plates were used for thin layer chromatography. Both flash and gravity chromatography were carried out on E. Merck Kieselgel 60 (230-400 mesh) silica gel. Analytical and preparative HPLC were carried out on Rainin or Beckman chromatographs. ODS refers to an octadecylsilyl derivatized silica gel chromatographic support. 5 μ Apex-ODS indicates an octadecylsilyl derivatized silica gel chromatographic support having a nominal particle size of 5 μ , made by Jones Chromatography, Littleton, Colorado. YMC ODS-AQ® is an ODS chromatographic support and is a registered trademark of YMC Co. Ltd., Kyoto, Japan. PRP-1® is a polymeric (styrene-divinylbenzene) chromatographic support, and is a registered trademark of Hamilton Co., Reno, Nevada) Celite® is a filter aid composed of acid-washed diatomaceous silica, and is a registered trademark of Manville Corp., Denver, Colorado.

All reagents and solvents were obtained from commercial vendors. Starting materials (e.g., amines and epoxides) were synthesized using standard techniques and procedures.

With appropriate manipulation and protection of any chemical functionality, synthesis of the remaining compounds of Formula (I) is accomplished by methods analogous to those above and to those described in the Experimental section.

In order to use a compound of Formula (I) or a pharmaceutically acceptable salt thereof for the treatment of humans and other mammals, it is normally formulated in accordance with standard pharmaceutical practice as a pharmaceutical composition.

The calcilytic compounds can be administered by different routes including intravenous, intraperitoneal, subcutaneous, intramuscular, oral, topical (transdermal), or transmucosal administration. For systemic administration, oral administration is preferred. For oral administration, for example, the compounds can be formulated into conventional oral dosage forms such as capsules, tablets, and liquid preparations such as syrups, elixirs, and concentrated drops.

Alternatively, injection (parenteral administration) may be used, *e.g.*, intramuscular, intravenous, intraperitoneal, and subcutaneous. For injection, the compounds of the invention are formulated in liquid solutions, preferably, in physiologically compatible buffers or solutions, such as saline solution, Hank's solution, or Ringer's solution. In addition, the compounds may be formulated in solid form and redissolved or suspended immediately prior to use. Lyophilized forms can also be produced.

Systemic administration can also be by transmucosal or transdermal means. For transmucosal or transdermal administration, penetrants appropriate to the barrier to be permeated are used in the formulation. Such penetrants are generally known in the art, and include, for example, for transmucosal administration, bile salts and fusidic acid derivatives. In addition, detergents may be used to facilitate permeation. Transmucosal administration, for example, may be through nasal sprays, rectal suppositories, or vaginal suppositories.

For topical administration, the compounds of the invention can be formulated into ointments, salves, gels, or creams, as is generally known in the art.

The amounts of various calcilytic compounds to be administered can be determined by standard procedures taking into account factors such as the compound IC_{50} , EC_{50} , the biological half-life of the compound, the age, size and weight of the patient, and the disease or disorder associated with the patient. The importance of these and other factors to be considered are known to those of ordinary skill in the art.

Amounts administered also depend on the routes of administration and the degree of oral bioavailability. For example, for compounds with low oral bioavailability, relatively higher doses will have to be administered.

Preferably the composition is in unit dosage form. For oral application, for example, a tablet, or capsule may be administered, for nasal application, a metered aerosol dose may be administered, for transdermal application, a topical formulation or patch may be administered and for transmucosal delivery, a buccal patch may be administered. In each case, dosing is such that the patient may administer a single dose.

Each dosage unit for oral administration contains suitably from 0.01 to 500 mg/Kg, and preferably from 0.1 to 50 mg/Kg, of a compound of Formula (I) or a

pharmaceutically acceptable salt thereof, calculated as the free base. The daily dosage for parenteral, nasal, oral inhalation, transmucosal or transdermal routes contains suitably from 0.01 mg to 100 mg/Kg, of a compound of Formula(I). A topical formulation contains suitably 0.01 to 5.0% of a compound of Formula (I).

- 5 The active ingredient may be administered, for example, from 1 to 6 times per day, preferably once, sufficient to exhibit the desired activity, as is readily apparent to one skilled in the art.

As used herein, "treatment" of a disease includes, but is not limited to prevention, retardation and prophylaxis of the disease.

- 10 Diseases and disorders which might be treated or prevented, based upon the affected cells, include bone and mineral-related diseases or disorders; hypoparathyroidism; those of the central nervous system such as seizures, stroke, head trauma, spinal cord injury, hypoxia-induced nerve cell damage, such as occurs in cardiac arrest or neonatal distress, epilepsy, neurodegenerative diseases such as
- 15 Alzheimer's disease, Huntington's disease and Parkinson's disease, dementia, muscle tension, depression, anxiety, panic disorder, obsessive-compulsive disorder, post-traumatic stress disorder, schizophrenia, neuroleptic malignant syndrome, and Tourette's syndrome; diseases involving excess water reabsorption by the kidney, such as syndrome of inappropriate ADH secretion (SIADH), cirrhosis, congestive
- 20 heart failure, and nephrosis; hypertension; preventing and/or decreasing renal toxicity from cationic antibiotics (*e.g.*, aminoglycoside antibiotics); gut motility disorders such as diarrhea and spastic colon; GI ulcer diseases; GI diseases with excessive calcium absorption such as sarcoidosis; autoimmune diseases and organ transplant rejection; squamous cell carcinoma; and pancreatitis.

- 25 In a preferred embodiment of the present invention, the present compounds are used to increase serum parathyroid hormone ("PTH") levels. Increasing serum PTH levels can be helpful in treating diseases such as hypoparathyroidism, osteosarcoma, periodontal disease, fracture, osteoarthritis, rheumatoid arthritis, Paget's disease, humoral hypercalcemia malignancy and osteoporosis.

- 30 Another aspect of the present invention describes a method of treating a patient comprising administering to the patient an amount of a present compound sufficient to increase the serum PTH level. Preferably, the method is carried out by

administering an amount of the compound effective to cause an increase in duration and/or quantity of serum PTH level sufficient to have a therapeutic effect.

In various embodiments, the compound administered to a patient causes an increase in serum PTH having a duration of up to one hour, about one to about
5 twenty-four hours, about one to about twelve hours, about one to about six hours, about one to about five hours, about one to about four hours, about two to about five hours, about two to about four hours, or about three to about six hours.

In an alternate embodiment of the present invention, the compound administered causes an increase in serum PTH of longer than about twenty-four
10 hours, but the compound is co-administered with an anti-resorptive agent.

In additional different embodiments, the compound administered to a patient causes an increase in serum PTH of up to two fold, two to five fold, five to ten fold, and at least 10 fold, greater than peak serum PTH in the patient. The peak serum level is measured with respect to a patient not undergoing treatment.

15 Composition of Formula (I) and their pharmaceutically acceptable salts, which are active when given orally, can be formulated as syrups, tablets, capsules and lozenges. A syrup formulation will generally consist of a suspension or solution of the compound or salt in a liquid carrier for example, ethanol, peanut oil, olive oil, glycerine or water with a flavoring or coloring agent. Where the
20 composition is in the form of a tablet, any pharmaceutical carrier routinely used for preparing solid formulations may be used. Examples of such carriers include magnesium stearate, terra alba, talc, gelatin, acacia, stearic acid, starch, lactose and sucrose. Where the composition is in the form of a capsule, any routine encapsulation is suitable, for example using the aforementioned carriers in a hard
25 gelatin capsule shell. Where the composition is in the form of a soft gelatin shell capsule any pharmaceutical carrier routinely used for preparing dispersions or suspensions may be considered, for example aqueous gums, celluloses, silicates or oils, and are incorporated in a soft gelatin capsule shell.

Typical parenteral compositions consist of a solution or suspension of a
30 compound or salt in a sterile aqueous or non-aqueous carrier optionally containing a parenterally acceptable oil, for example polyethylene glycol, polyvinylpyrrolidone, lecithin, arachis oil or sesame oil.

Typical compositions for inhalation are in the form of a solution, suspension or emulsion that may be administered as a dry powder or in the form of an aerosol using a conventional propellant such as dichlorodifluoromethane or trichlorofluoromethane.

5 A typical suppository formulation comprises a compound of Formula (I) or a pharmaceutically acceptable salt thereof which is active when administered in this way, with a binding and/or lubricating agent, for example polymeric glycols, gelatins, cocoa-butter or other low melting vegetable waxes or fats or their synthetic analogs.

10 Typical dermal and transdermal formulations comprise a conventional aqueous or non-aqueous vehicle, for example a cream, ointment, lotion or paste or are in the form of a medicated plaster, patch or membrane.

Preferably the composition is in unit dosage form, for example a tablet, capsule or metered aerosol dose, so that the patient may administer a single dose.

15 No unacceptable toxicological effects are expected when compounds of the present invention are administered in accordance with the present invention.

The biological activity of the compounds of Formula (I) are demonstrated by the following tests:

(I) Calcium Receptor Inhibitor Assay

20 Calcilytic activity was measured by determining the IC_{50} of the test compound for blocking increases of intracellular Ca^{2+} elicited by extracellular Ca^{2+} in HEK 293 4.0-7 cells stably expressing the human calcium receptor. HEK 293 4.0-7 cells were constructed as described by Rogers *et al.*, *J. Bone Miner. Res.* 10 Suppl. 1:S483, 1995 (hereby incorporated by reference herein). Intracellular Ca^{2+} increases were elicited by increasing extracellular Ca^{2+} from 1 to 1.75 mM. Intracellular Ca^{2+} was measured using fluo-3, a fluorescent calcium indicator.

The procedure was as follows:

1. Cells were maintained in T-150 flasks in selection media (DMEM supplemented with 10% fetal bovine serum and 200 ug/mL hygromycin B), under
30 5% CO_2 :95% air at 37 °C and were grown up to 90% confluency.

2. The medium was decanted and the cell monolayer was washed twice with phosphate-buffered saline (PBS) kept at 37 °C. After the second wash, 6 mL

of 0.02% EDTA in PBS was added and incubated for 4 minutes at 37 °C.

Following the incubation, cells were dispersed by gentle agitation.

3. Cells from 2 or 3 flasks were pooled and pelleted (100 x g). The cellular pellet was resuspended in 10-15 mL of SPF-PCB+ and pelleted again by centrifugation. This washing was done twice.

Sulfate- and phosphate-free parathyroid cell buffer (SPF-PCB) contains 20 mM Na-Hepes, pH 7.4, 126 mM NaCl, 5 mM KCl, and 1 mM MgCl₂. SPF-PCB was made up and stored at 4 °C. On the day of use, SPF-PCB was supplemented with 1 mg/mL of D-glucose and 1 mM CaCl₂ and then split into two fractions. To one fraction, bovine serum albumin (BSA; fraction V, ICN) was added at 5 mg/mL (SPF-PCB+). This buffer was used for washing, loading and maintaining the cells. The BSA-free fraction was used for diluting the cells in the cuvette for measurements of fluorescence.

4. The pellet was resuspended in 10 mL of SPF-PCB+ containing 2.2 μM fluo-3 (Molecular Probes) and incubated at room temperature for 35 minutes.

5. Following the incubation period, the cells were pelleted by centrifugation. The resulting pellet was washed with SPF-PCB+. After this washing, cells were resuspended in SPF-PCB+ at a density of 1-2 x 10⁶ cells/mL.

6. For recording fluorescent signals, 300 μL of cell suspension were diluted in 1.2 mL of SPF buffer containing 1 mM CaCl₂ and 1 mg/mL of D-glucose. Measurements of fluorescence were performed at 37 °C with constant stirring using a spectrofluorimeter. Excitation and emission wavelengths were measured at 485 and 535 nm, respectively. To calibrate fluorescence signals, digitonin (5 mg/mL in ethanol) was added to obtain F_{max}, and the apparent F_{min} was determined by adding Tris-EGTA (2.5 M Tris-Base, 0.3 M EGTA). The concentration of intracellular calcium was calculated using the following equation: Intracellular calcium = $(F - F_{\min} / F_{\max}) \times K_d$; where K_d = 400 nM.

7. To determine the potential calcilytic activity of test compounds, cells were incubated with test compound (or vehicle as a control) for 90 seconds before increasing the concentration of extracellular Ca²⁺ from 1 to 2 mM. Calcilytic compounds were detected by their ability to block, in a concentration-dependent manner, increases in the concentration of intracellular Ca²⁺ elicited by extracellular Ca²⁺.

In general, those compounds having lower IC₅₀ values in the Calcium Receptor Inhibitor Assay are more preferred compounds. Compounds having an IC₅₀ greater than 50 uM were considered to be inactive. Preferred compounds are those having an IC₅₀ of 10uM or lower, more preferred compounds have an IC₅₀ of 1uM, and most preferred compounds have an IC₅₀ of 0.1uM or lower.

(II) Calcium Receptor Binding Assay

HEK 293 4.0-7 cells stably transfected with the Human Parathyroid Calcium Receptor("HuPCaR") were scaled up in T180 tissue culture flasks. Plasma membrane is obtained by polytron homogenization or glass douncing in buffer (50mM Tris-HCl pH 7.4, 1mM EDTA, 3mM MgCl₂) in the presence of a protease inhibitor cocktail containing 1uM Leupeptin, 0.04 uM Pepstatin, and 1 mM PMSF. Aliquoted membrane was snap frozen and stored at -80°C. ³H labeled compound was radiolabeled to a radiospecific activity of 44Ci/mmol and was aliquoted and stored in liquid nitrogen for radiochemical stability.

A typical reaction mixture contains 2 nM ³H compound ((R,R)-N-4'-Methoxy-t-3-3'-methyl-1'-ethylphenyl-1-(1-naphthyl)ethylamine), or ³H compound (R)-N-[2-Hydroxy-3-(3-chloro-2-cyanophenoxy)propyl]-1,1-dimethyl-2-(4-methoxyphenyl)ethylamine 4-10 ug membrane in homogenization buffer containing 0.1% gelatin and 10% EtOH in a reaction volume of 0.5 mL. Incubation is performed in 12 x 75 polyethylene tubes in an ice water bath. To each tube 25 uL of test sample in 100% EtOH is added, followed by 400 uL of cold incubation buffer, and 25 uL of 40 nM ³H-compound in 100% EtOH for a final concentration of 2nM. The binding reaction is initiated by the addition of 50 uL of 80-200 ug/mL HEK 293 4.0-7 membrane diluted in incubation buffer, and allowed to incubate at 4°C for 30 min. Wash buffer is 50 mM Tris-HCl containing 0.1% PEI. Nonspecific binding is determined by the addition of 100-fold excess of unlabeled homologous ligand, and is generally 20% of total binding. The binding reaction is terminated by rapid filtration onto 1% PEI pretreated GF/C filters using a Brandel Harvester. Filters are placed in scintillation fluid and radioactivity assessed by liquid scintillation counting.

The following examples are illustrative, but not limiting of the embodiments of the present invention.

Example 1

5 Preparation of (R)-N-[2-Hydroxy-3-(3-chloro-2-cyanophenoxy)propyl]-1,1-
 dimethyl-2-(2,3-dihydrobenzo[b]furan-5yl)ethylamine Hydrochloride
 5-(2-Amino-2-methylpropyl)-2,3-dihydrobenzo[b]furan

 Sodium hydride (0.89g, 37.1 mmole) was added to 45 mL of DMSO and
10 stirred for 30 min at room temperature. Isopropyltriphenylphosphonium iodide
 (16.05g, 37.1 mmole) was then added and stirred for 1.5 hours followed by the
 addition of 2,3-dihydrobenzo[b]furan-5-carboxaldehyde (5.0g, 33.75 mmole). This
 mixture was stirred for 18 hours at room temperature then poured into water (300
 mL) / conc. HCl (5 mL) and extracted with ether. The crude product was
15 chromatographed on silica gel in 1% MeOH/CHCl₃ to afford 5.1 g (87%) of 5-(2-
 methylpropenyl)-2,3-dihydrobenzo[b]furan, which was 100% pure by GC-MS.
 To a 0°C suspension of sodium cyanide (1.44g, 29.3 mmole) in 6 mL of acetic acid
 was slowly added a 0°C solution of sulfuric acid (3.2 mL) in acetic acid (3.2 mL).
 After stirring for 45 min at 0°C, 5-(2-methylpropenyl)-2,3-dihydrobenzo[b]furan
20 (5.1g, 29.3 mmole) was added, and the mixture allowed to warm to room
 temperature while stirring for 18 hours. The reaction was poured into ice/ NaOH
 and extracted with ether. The ether layer was dried over sodium sulfate, then
 concentrated in vacuo. The crude amidated product was taken up in EtOH/NaOH
 and refluxed for 24 hours. The ethanol was removed in vacuo, and the residue
25 taken up in ether and water. The ether layer was separated, dried over sodium
 sulfate and concentrated in vacuo to yield the crude amine as a dark oil. The
 product was purified by short-path distillation at reduced pressure.
 (R)-N-[2-Hydroxy-3-(3-chloro-2-cyanophenoxy)propyl]-1,1-dimethyl-2-(2,3-
 dihydrobenzo[b]furan-5yl)ethylamine Hydrochloride

30 Using previously described methods, (R)-3-chloro-2-cyanophenyl glycidyl
 ether (0.398 g, 1.9 mmol) and 5-(2-amino-2-methylpropyl)-2,3-
 dihydrobenzo[b]furan (0.382 g, 2.0 mmol) were used to prepare 100 mg of the title
 compound as a white solid. ¹H-NMR (CDCl₃) • 9.65 (1H, m), 8.13 (1H, m), 7.4

(1H, t), 7.05 (2H, d), 6.92 (2H, d), 6.65 (1H, d), 5.7 (1H, d), 4.77 (1H, br m), 4.53 (2H, t), 4.25 (2H, d), 3.4 (2H, m), 3.1 (4H, m), 1.4 (6H, d).

Example 2

5

Preparation of (R)-N-[2-Hydroxy-3-(3-chloro-2-cyanophenoxy)propyl]-1,1-dimethyl-2-(quinolin-3-yl)ethylamine Dihydrochloride
3-(2-Amino-2-methylpropyl)quinoline

10 To a solution of 3-quinolinecarboxaldehyde (10.27g, 65.3 mmole) in 10 mL of pyridine and 30 mL of ethanol was added methoxylamine hydrochloride (6.0 g, 71.9 mmole). After stirring for 2 hours the solvents were removed under reduced pressure, and the residue taken up in ether and water. The ether layer was separated, dried over sodium sulfate and concentrated in vacuo. The crude oxime
15 (11.91 g, 63.9 mmole) was dissolved in 120 mL of trifluoroacetic acid and treated with zinc powder (13.0 g, 199 mmole). After 10 min, the reaction refluxed spontaneously for a few seconds, and the mixture was stirred for another 3 hours. The mixture was poured into water, and washed with ether. The aqueous layer was then made basic with NaOH, and the amine extracted into ether. The ether layer
20 was separated, dried over sodium sulfate and concentrated in vacuo to yield 8.68 g of 3-(aminomethyl)quinoline. To this amine (8.68 g, 54.9 mmole), dissolved in 200 mL of dichloromethane, was added 2,4,6-triphenylpyrylium tetrafluoroborate (19.56 g, 49.4 mmole), and the reaction stirred at room temperature for 48 hours. The solids were filtered off, and the resulting solution concentrated in vacuo to give
25 25.4 g (86.3%) of the crude N-(3-quinolinylmethyl)-2,4,6-triphenylpyridinium tetrafluoroborate salt. A solution of this salt (25.4 g, 47.4 mmole) in 100 mL of DMSO was added to the sodium salt of 2-nitropropane (142.1 mmole) (made by adding sodium hydride (3.41 g, 142 mmole) to 50 mL of methanol followed by addition of 2-nitropropane (12.66 g, 142.1 mmole), then removing the methanol in
30 vacuo). The reaction was stirred for 24 hours at 100 C then cooled and diluted with ether and aqueous HCl. The aqueous layer was separated, made basic with NaOH, and extracted with ether. The ether layer was dried over sodium sulfate, and concentrated in vacuo to give after purification on silica gel (in chloroform) 10.7 g

(98%) of 3-(2-nitro-2-methylpropyl)quinoline. To this nitro compound (10.7 g, 47.2 mmole) dissolved in 100 mL of trifluoroacetic acid was slowly added zinc powder (9.3 g, 142 mmole). Stirred for 24 hours at room temperature. The reaction mixture was then poured into water and washed with ether. The aqueous layer was separated, made basic with NaOH, and extracted with ether. The ether layer was dried over sodium sulfate, and concentrated in vacuo to give 4.5 g (48%) of 3-(2-amino-2-methylpropyl)quinoline. GC/EI-MS, m/z , (rel. int.) 185 (M^+ - 15, 3), 143 (58), 115 (11), 58 (100), 42 (7).

10 (R)-N-[2-Hydroxy-3-(3-chloro-2-cyanophenoxy)propyl]-1,1-dimethyl-2-(quinolin-3-yl)ethylamine Dihydrochloride

Using previously described methods, (R)-3-chloro-2-cyanophenyl glycidyl ether (0.398g, 1.9 mmol) and 3-(2-amino-2-methylpropyl)quinoline (0.401 g, 2.0 mmol) were used to prepare 130 mg of the title compound as a white solid. ^1H -NMR

15 (CDCl_3) • 9.7 (1H, br t), 9.25 (1H, s), 9.0 (2H, br s), 8.42 (1H, d), 8.37 (1H, d), 8.1 (1H, dd), 7.93 (1H, dd), 7.65 (1H, dd), 7.35 (1H, d), 7.28 (1H, d), 4.3 (3H, m), 3.45 (2H, s), 3.3 (2H, m), 2.5 (1H, s), 1.4 (6H, s).

Example 3

20

Preparation of (R)-N-[2-Hydroxy-3-(3-chloro-2-cyanophenoxy)propyl]-1,1-dimethyl-2-(quinolin-2-yl)ethylamine Dihydrochloride
2-(2-Amino-2-methylpropyl)quinoline

25

Using the method of Example 2, *supra*, 2-(2-amino-2-methylpropyl)quinoline was prepared from quinoline-2-carboxaldehyde. GC/EI-MS, m/z , (rel. int.) 185 (M^+ - 15, 5), 143 (42), 115 (13), 58 (100), 42 (6). (R)-N-[2-Hydroxy-3-(3-chloro-2-cyanophenoxy)propyl]-1,1-dimethyl-2-(quinolin-2-yl)ethylamine Dihydrochloride

30

Using previously described methods (R)-3-chloro-2-cyanophenyl glycidyl ether (0.21g, 1.0 mmol) and 2-(2-amino-2-methylpropyl)quinoline (0.24 g, 1.2 mmol) were used to prepare 24 mg of the title compound as a white solid. ^1H -NMR (CDCl_3) • 9.7 (1H, m), 9.3 (1H, m), 8.55 (1H, d), 8.39 (1H, d), 7.83 (1H, d), 7.7

(2H, m), 7.53 (1H, m), 7.15 (1H, t), 6.75 (2H, m), 4.3 (1H, m), 4.0 (2H, m), 3.65 (2H, dd), 3.15 (2H, m), 1.3 (6H, d).

Example 4

5

Preparation of (R)-N-[2-Hydroxy-3-(3-chloro-2-cyanophenoxy)propyl]-1,1-dimethyl-2-(isoquinolin-3-yl)ethylamine Dihydrochloride
3-(2-Amino-2-methylpropyl)isoquinoline

10 Using the method of Example 2, *supra*, 3-(2-amino-2-methylpropyl)isoquinoline was prepared from isoquinoline-3-carboxaldehyde. GC/EI-MS, *m/z*, (rel. int.) 185 (*M*⁺ - 15, 7), 144 (13), 143 (100), 116 (9), 115 (22), 58 (47), 42 (8).

(R)-N-[2-Hydroxy-3-(3-chloro-2-cyanophenoxy)propyl]-1,1-dimethyl-2-
15 (isoquinolin-3-yl)ethylamine Dihydrochloride

Using previously described methods (R)-3-chloro-2-cyanophenyl glycidyl ether (0.47g, 2.24 mmol) and 3-(2-amino-2-methylpropyl)isoquinoline (0.49 g, 2.45 mmol) were used to prepare 200 mg of the title compound as a light yellow solid. ¹H-NMR (CDCl₃) • 9.7 (2H, s on top of m), 9.25 (1H, m), 8.45 (1H, d), 8.27 (1H, s), 8.2 (1H, d), 8.1 (1H, t), 7.9 (1H, t), 7.68 (1H, t), 7.35 (1H, d), 7.25 (1H, d), 4.39
20 (1H, m), 4.3 (2H, s), 3.57 (2H, dd), 3.3 (2H, m), 1.4 (6H, d).

Example 5

25 Preparation of (R)-N-[2-Hydroxy-3-(3-chloro-2-cyanophenoxy)propyl]-1,1-dimethyl-4-(2-pyridyl)butylamine Dihydrochloride
4-Benzylloxycarbonylamino-4-methylpent-1-ene

The 2,2-dimethyl-4-pentenoic acid (20.7g, 162 mmoles) was dissolved in
30 300 mL of benzyl alcohol followed by addition of triethylamine (17.98g, 178 mmoles). Diphenyl phosphorylazide (46.67g, 170 mmoles) was added and the reaction heated to 100°C overnight under nitrogen. The product was separated from

the excess benzyl alcohol by distillation. The product distilled at approx. 130°C @ 0.01 mm.

9-(4-Benzyloxycarbonylamino-4-methylpentyl)-9-borabicyclo[3.3.1]nonane

To a 0.5 M solution of 9-BBN in THF (100 mL, 50 mmole) was added 4-benzyloxycarbonylamino-4-methylpent-1-ene (11.67 g, 50 mmole). The reaction was allowed to stand for 24 hours at room temperature. Analysis by GC-MS showed no starting alkene left. The solution, which was approximately 0.439 M in the borane, was used without purification.

2-(4-Amino-4-methylpentyl)pyridine

To 2-bromopyridine (0.948 g, 6 mmole) was added 9-(4-benzyloxycarbonylamino-4-methylpentyl)-9-borabicyclo[3.3.1]nonane (12 mL, 5 mmole of a THF solution) in a nitrogen flushed reaction tube. To this solution was added 0.122 g (0.15 mmole) of [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II) dichloromethane complex (1:1), 1.38 g (10 mmole) of potassium carbonate, and 1.25 mL of water. The reaction was stirred for 18 hours at 65 C, then poured into aqueous NaOH, and extracted with ether. The ether layer was separated, washed with brine, dried over sodium sulfate, and concentrated in vacuo. The crude product was taken up in 20 mL of ethanol to which 1 g of palladium hydroxide on carbon (10%) was added. The mixture was stirred for 18 hours under a hydrogen balloon. The reaction mixture was filtered and concentrated in vacuo. The residue was taken up in aqueous HCl, and extracted with ether. The aqueous layer was separated, made basic with NaOH, and extracted with ether. The ether layer was dried over sodium sulfate, and concentrated in vacuo to give 0.63 g of 2-(4-Amino-4-methylpentyl)pyridine.

(R)-N-[2-Hydroxy-3-(3-chloro-2-cyanophenoxy)propyl]-1,1-dimethyl-4-(2-pyridyl)butylamine Dihydrochloride

Using previously described methods (R)-3-chloro-2-cyanophenyl glycidyl ether (0.21g, 1.0 mmol) and 2-(4-Amino-4-methylpentyl)pyridine (0.196 g, 1.1 mmol) were used to prepare 19 mg of the title compound as a light yellow solid. LC/MS/MS* (In-Source ISD technique), m/z , 388 (M^+), 227, 162, 106.

Example 6

Preparation of (R)-N-[2-Hydroxy-3-(3-chloro-2-cyano-4-
morpholinosulfonamidophenoxy)propyl]-1,1-dimethyl-4-(2-pyridyl)butylamine
5 Dihydrochloride

Using the method of Example 5, *supra*, (R)-3-chloro-2-cyano-4-
morpholinosulfonamidophenyl glycidyl ether (0.3g, 0.83 mmol) and 2-(4-Amino-4-
methylpentyl)pyridine (0.156 g, 0.87 mmol) were used to prepare 150 mg of the
10 title compound as an off white solid. ¹H-NMR (CDCl₃) δ 9.3 (1H, m), 8.8 (2H, d
on top of m), 8.55 (1H, t), 8.18 (1H, d), 8.02 (1H, d), 7.95 (1H, dd), 7.5 (1H, d), 4.4
(3H, m), 3.6 (4H, br s), 3.1 (6H, m), 2.5 (2H, s), 1.8 (4H, m), 1.3 (6H, s).

Example 7

Preparation of (R)-N-[2-Hydroxy-3-(3-chloro-2-cyanophenoxy)propyl]-1,1-
dimethyl-4-(3-pyridyl)butylamine Dihydrochloride
20 3-(4-Amino-4-methylpentyl)pyridine

Using the method of Example 5, *supra*, 0.66 g of 3-(4-Amino-4-
methylpentyl)pyridine was prepared, starting with 6 mmoles of 3-bromopyridine.
(R)-N-[2-Hydroxy-3-(3-chloro-2-cyanophenoxy)propyl]-1,1-dimethyl-4-(3-
pyridyl)butylamine Dihydrochloride

25 Using previously described methods, (R)-3-chloro-2-cyanophenyl glycidyl
ether (0.21g, 1.0 mmol) and 3-(4-Amino-4-methylpentyl)pyridine (0.196 g, 1.1
mmol) were used to prepare 25 mg of the title compound as a light yellow glassy
solid. LC/MS/MS* (In-Source ISD technique), *m/z*, 388 (M⁺), 227, 162, 106.

Example 8

Preparation of (R)-N-[2-Hydroxy-3-(3-chloro-2-cyano-4-
morpholinosulfonamidophenoxy)propyl]-1,1-dimethyl-4-(3-pyridyl)butylamine
5 Dihydrochloride

Using the method of Example 5, *supra*, (R)-3-chloro-2-cyano-4-morpholinosulfonamidophenyl glycidyl ether (0.3g, 0.83 mmol) and 3-(4-Amino-4-methylpentyl)pyridine (0.156 g, 0.87 mmol) were used to prepare 30 mg of the
10 title compound as an off white solid. ¹H-NMR (CDCl₃) δ 9.3 (1H, m), 8.9 (1H, s), 8.8 (2H, d on top of m), 8.56 (1H, d), 8.16 (1H, d), 8.04 (1H, dd), 7.5 (1H, d), 4.4 (3H, m), 3.6 (4H, br s), 3.1 (4H, m), 2.8 (2H, br s), 2.5 (2H, s), 1.7 (4H, m), 1.3 (6H, s).

15

Example 9

Preparation of (R)-N-[2-Hydroxy-3-(3-chloro-2-cyanophenoxy)propyl]-4-
phenylbutylamine Hydrochloride

20 Using previously described methods, (R)-3-chloro-2-cyanophenyl glycidyl ether (0.21g, 1.0 mmol) and 4-phenylbutylamine (0.164 g, 1.1 mmol) were used to prepare 250 mg of the title compound as a white solid. ¹H NMR (CDCl₃) δ 10.09 (1H, s), 9.38 (1H, s), 9.12 (1H, s), 7.53 (2H, m), 7.19 (7H, m), 6.08 (1H, m), 5.98 (1H, m), 4.63 (1H, m), 4.52 (1H, m), 4.23 (3H, m), 3.95 (1H, m), 3.43 (1H, m), 3.20
25 (1H, m), 3.00 (1H, m), 2.65 (3H, m), 1.86 (2H, m), 1.71 (2H, m).

Example 10

Preparation of (R)-N-[2-Hydroxy-3-(3-chloro-2-cyanophenoxy)propyl]-1,1-
30 dimethyl-4-(2-carbethoxyphenyl)butylamine Hydrochloride
Ethyl 2-(4-Amino-4-methylpentyl)benzoate

To ethyl 2-bromobenzoate (0.504 g, 2.2 mmole) in a nitrogen flushed reaction tube was added 0.049 g (0.06 mmole) of [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II) dichloromethane complex (1:1) dissolved in 2 mL of DMF. To this solution was added 1.3 g (4 mmole) of cesium carbonate, followed by 9-(4-benzyloxycarbonylamino-4-methylpentyl)-9-borabicyclo[3.3.1]nonane (4.56 mL, 2.0 mmole of a THF solution). The reaction was stirred for 16.5 hours at 50 C, then poured into aqueous NaOH, and extracted with ether. The ether layer was separated, washed with brine, dried over sodium sulfate, and concentrated in vacuo. The crude product was taken up in 10 mL of ethanol to which 0.3 g of palladium hydroxide on carbon (10%) was added. The mixture was stirred for 18 hours under a hydrogen balloon. The reaction mixture was filtered and concentrated in vacuo. The residue was taken up in aqueous HCl, and extracted with ether. The aqueous layer was separated, made basic with NaOH, and extracted with ether. The ether layer was dried over sodium sulfate, and concentrated in vacuo to give crude ethyl 2-(4-amino-4-methylpentyl)benzoate. The crude product was purified by reversed-phase HPLC on a C-18 column using a gradient of 0.1% HCl to 40% acetonitrile in 0.1% HCl.

(R)-N-[2-Hydroxy-3-(3-chloro-2-cyanophenoxy)propyl]-1,1-dimethyl-4-(2-carbethoxyphenyl)butylamine Hydrochloride

Using previously described methods, (R)-3-chloro-2-cyanophenyl glycidyl ether (0.21g, 1.0 mmol) and 1,1-dimethyl-4-(2-carbethoxyphenyl)butylamine (0.274 g, 1.1 mmol) were used to prepare 260 mg of the title compound as a white solid. ¹H NMR (CDCl₃) • 9.54 (1H, s), 8.17 (1H, m), 7.85 (1H, dd), 7.43 (2H, m), 7.24 (2H, m), 7.06 (1H, d), 6.97 (1H, d), 6.00 (1H, d), 4.71 (1H, s), 4.33 (2H, q), 4.26 (2H, d), 3.27 (2H, m), 2.60 (2H, m), 1.85 (2H, m), 1.69 (2H, m), 1.46 (6H, s), 1.37 (3H, t).

Example 11

Preparation of (R)-N-[2-Hydroxy-3-(3-chloro-2-cyanophenoxy)propyl]-1,1-dimethyl-4-(3-carbethoxyphenyl)butylamine Hydrochloride

Using previously described methods, (R)-3-chloro-2-cyanophenyl glycidyl ether (0.21g, 1.0 mmol) and 1,1-dimethyl-4-(3-carbethoxyphenyl)butylamine (0.274 g, 1.1 mmol) were used to prepare 230 mg of the title compound as a white solid. ¹H NMR (CDCl₃) • 9.56 (1H, m), 8.21 (1H, m), 7.86 (2H, m), 7.37 (3H, m), 7.07 (1H, d), 6.94 (1H, d), 5.59 (1H, d), 4.70 (1H, m), 4.35 (2H, q), 4.24 (2H, d), 3.25 (2H, m), 2.71 (2H, m), 1.82 (4H, m), 1.49 (6H, s), 1.39 (3H, t).

Example 12

Preparation of (R)-N-[2-Hydroxy-3-(3-chloro-2-cyanophenoxy)propyl]-1,1-dimethyl-4-(4-carbethoxyphenyl)butylamine Hydrochloride

Using the method of Example 5, *supra*, (R)-3-chloro-2-cyanophenyl glycidyl ether (0.21g, 1.0 mmol) and 1,1-dimethyl-4-(4-carbethoxyphenyl)butylamine (0.274 g, 1.1 mmol) were used to prepare 250 mg of the title compound as a white solid. ¹H NMR (CDCl₃) • 9.56 (1H, m), 8.19 (1H, m), 7.95 (2H, d), 7.43 (1H, ddd), 7.26 (2H, d), 7.07 (1H, d), 6.93 (1H, d), 5.58 (1H, d), 4.69 (1H, m), 4.33 (2H, q), 4.22 (2H, d), 3.23 (2H, m), 2.71 (2H, m), 1.48 (3H, s), 1.47 (3H, s), 1.37 (3H, t).

Example 13

Preparation of (R)-N-[2-Hydroxy-3-(3-chloro-2-cyanophenoxy)propyl]-1,1-dimethyl-2-(4-ethylpyrid-2-yl)ethylamine Dihydrochloride 1,1-Dimethyl-2-(4-ethylpyrid-2-yl)ethylamine

4 mmoles 5-ethyl-2-methyl pyridine in 4 mL dry ether was treated with 4.32 mmoles of phenyl lithium (1.8 M solution in cyclohexane/ether) at 0°C. After reaction at RT for 1 h the solution was added dropwise to a chilled (ice bath) solution of 2 mmoles isopropylidene-3-nitrobenzene sulfenamide in 2 mL dry ether. After reaction at RT for 1 hr and at reflux for 0.5 hrs the cooled reaction mixture was quenched with 5 mls water. The organic layer was extracted three

times with 6 Molar HCl. The pooled HCl extracts were evaporated to an oil, made basic with 10 N NaOH and extracted with ether. The ether was extracted twice with pH 7 Phosphate Buffer, buffer extracts made basic with NaOH and extracted with chloroform. Removal of the chloroform resulted in the title compound in 24%

5 yield.

(R)-N-[2-Hydroxy-3-(3-chloro-2-cyanophenoxy)propyl]-1,1-dimethyl-2-(4-ethylpyrid-2-yl)ethylamine Dihydrochloride

Using previously described methods, (R)-3-chloro-2-cyanophenyl glycidyl ether (0.21g, 1.0 mmol) and 1,1-Dimethyl-2-(4-ethylpyrid-2-yl)ethylamine (0.25 g,
10 1.4 mmol) were used to prepare 314 mg of the title compound as a white solid. ¹H-NMR (CDCl₃) • 9.95 (1H, br s), 9.05 (1H, br s), 8.62 (1H,s), 8.27 (1H, br s), 8.02 (1H, br s), 7.44 (1H, t), 7.0 (2H, d), 5.6 (2H, br s), 4.7 (1H, m), 4.34 (2H, br s), 3.9 (2H, br s), 3.53 (2H, br s), 2.86 (2H, q), 1.62 (6H, s), 1.32 (3H, t).

15

Example 14

(R)-N-[2-Hydroxy-3-(3-chloro-2-cyanophenoxy)propyl]-1,1-dimethyl-2-benzamidoethylamine Hydrochloride

Using previously described methods, (R)-3-chloro-2-cyanophenyl glycidyl ether (0.00g, 00 mmol) and 1,1-Dimethyl-2-benzamidoethylamine (00 g, 00 mmol) were used to prepare 000 mg of the title compound as a white solid. ¹H-NMR (CDCl₃) • 9.52 (1H, m), 8.40 (1H, m), 8.20 (1H, s) 7.95 (1H, d), 7.36 (4H, m), 7.02 (1H, d), 6.82 (1H, d), 4.70 (1H, m), 4.16 (2H, m), 3.82 (3H, m), 3.37 (2H, m), 1.51
25 (3H, s), 1.46 (3H, s); ¹³C NMR (CDCl₃) ••169.2, 161.3, 137.7, 134.7, 133.1, 131.9, 128.5, 127.8, 122.3, 113.8, 110.8, 103.2, 77.2, 70.8, 65.5, 61.4, 46.4, 44.8, 22.0, 21.6.

Example 15

30 (R)-N-[2-Hydroxy-3-(3-chloro-2-cyanophenoxy)propyl]-1,1-dimethyl-4-phenylbut-2-ynylamine Hydrochloride

Using previously described methods, (R)-3-chloro-2-cyanophenyl glycidyl ether (0.00g, 00 mmol) and 1,1-Dimethyl-4-phenylbut-2-ynylamine (00 g, 00 mmol) were used to prepare 000 mg of the title compound as a white solid. ¹H-NMR (CDCl₃) δ 9.97 (1H, m), 8.76 (1H, m), 7.41 (1H, ddd), 7.30 (3H, m), 7.18 (2H, m), 7.04 (1H, d), 6.92 (1H, d), 5.63 (1H, m), 4.76 (1H, m), 4.23 (2H, m), 3.51 (2H, m), 1.85 (3H, s), 1.83 (3H, s); ¹³C NMR (CDCl₃) δ 161.4, 137.8, 135.7, 134.5, 128.2, 127.9, 126.8, 122.2, 113.5, 110.9, 103.4, 86.8, 79.0, 70.9, 65.3, 55.5, 46.8, 26.8, 26.7, 24.7.

Example 16

10

Preparation of 1,1-dimethyl-2-[(ethyl-4-oxyacetate)-phenyl]ethylamine

A mixture of 1-nitro-1,1-dimethyl-2-(4-hydroxyphenyl)ethane (3.9g, 0.02mole), K₂CO₃ (2.76g, 0.02mole) and ethylbromoacetate (3.06g, 0.02mole) were refluxed in 75ml of acetone for 18h. The reaction was cooled to room temperature and filtered. The filtrate was concentrated in vacuo to yield 5.5g of an oil. This oil was dissolved in 75 ml of EtOH and 1 large spatula-full of washed Raney-nickel was added under argon. The mixture was hydrogenated at room temperature and 55 psi for 18 H. The reaction was filtered and the filtrate concentrated in vacuo to an oil which was filtered through a pad of silica gel eluting with 10% MeOH-CH₂Cl₂ (v/v). The first 200 ml were combined and concentrated in vacuo to yield 3.2g of a pale yellow oil. MS, m/z 252 (M+H), 503 (2M+H).

Formulations for pharmaceutical use incorporating compounds of the present invention can be prepared in various forms and with numerous excipients. Examples of such formulations are given below.

Example 17

Inhalant Formulation

30

A compound of Formula (I) (1 mg to 100 mg) is aerosolized from a metered dose inhaler to deliver the desired amount of drug per use.

Example 18
Tablet Formulation

	<u>Tablets/Ingredients</u>	<u>Per Tablet</u>
5	1. Active ingredient (Cmp. of Formula(I))	40 mg
	2. Corn Starch	20 mg
	3. Alginic acid	20 mg
	4. Sodium Alginate	20 mg
10	5. Mg stearate	13 mg

Procedure for tablet formulation

Ingredients 1, 2, 3 and 4 are blended in a suitable mixer/blender. Sufficient
 15 water is added portion-wise to the blend with careful mixing after each addition
 until the mass is of a consistency to permit its conversion to wet granules. The wet
 mass is converted to granules by passing it through an oscillating granulator using a
 No. 8 mesh (2.38 mm) screen. The wet granules are then dried in an oven at 140°F
 (60°C) until dry. The dry granules are lubricated with ingredient No. 5, and the
 20 lubricated granules are compressed on a suitable tablet press.

Example 19


Parenteral Formulation

25 A pharmaceutical composition for parenteral administration is prepared by
 dissolving an appropriate amount of a compound of Formula (I) in polyethylene
 glycol with heating. This solution is then diluted with water for injections (to 100
 ml). The solution is then rendered sterile by filtration through a 0.22 micron
 membrane filter and sealed in sterile containers.

30 All publications, including but not limited to patents and patent
 applications cited in this specification are herein incorporated by reference as if
 each individual publication were specifically and individually indicated to be
 incorporated by reference as though fully set forth.

What is claimed is:

1. A compound according to Formula (I) hereinbelow:



Formula (I)

5

wherein:

Y₁ is a covalent bond, alkylene or alkenylene of up to 4 carbon atoms, unsubstituted or substituted by C₁₋₄ alkyl or O;

Y₂ is methylene, unsubstituted or substituted by C₁₋₄ alkyl or haloalkyl;

- 10 Y₃ is covalent bond or selected from the group consisting of O, S, N-R^{IV}, C₁₋₄ alkylene-O, C₁₋₄ alkylene-S, and C₁₋₄ alkylene-N-R^{IV};

R^{IV} is selected from the group consisting of H, C₁₋₄ alkyl, and C₃₋₆ cycloalkyl;

R₃ and R₄ are, independently, methyl or ethyl, or, together, form cyclopropyl;

R₅ is heteroaryl or fused heteroaryl; wherein the hetero-ring contains N, O or S,

- 15 and is aromatic, dihydro or tetrahydro, unsubstituted or substituted with any substituents being selected from the group consisting of OH, OCH₃, CH(CH₃)₂, halogen, C₁₋₄ alkyl, C₁₋₄ alkoxy, C₃₋₆ cycloalkyl, OSO₂R^{IV}, CN, NO₂, OCF₃, CF₃, CH₂CF₃, (CH₂)_n CO₂H, (CH₂)_n CO₂R^{IV}, and O-(CH₂)_n CO₂R^{IV};
n is an integer from 0 to 3;

- 20 G is a covalent bond, CHR₆ or C-R₆, wherein R₆ is H, OH or O (forming a ketone);


R₇ is H, OH, or O-C₁₋₄ alkyl;

R₈ is H or C₁₋₄ alkyl; or R₇ and R₈ together form a ketone;


A and B are, independently, selected from the group consisting of a bond, CH₂,

- 25 NH, O, S and C=O, provided that either A or B is selected from CH₂ and NH; or A and B together form a bond; or the A-B moiety is represented by CH=CH or C≡C;


X is selected from sub formulas (Ia) to (Ie) hereinbelow:




(Ia)




(Ib)



(Ic)



(Id)



(1e)

wherein

- 5 W is selected from the group consisting of R_1 , SO_2R_1 , $C(O)R_1$, $SO_2NR_1R_1'$, $C(O)NR_1R_1'$, $C(O)OR_1$, and SO_3R_1' , wherein R_1 and R_1' are independently selected from the group consisting of hydrogen, C_{1-4} alkyl, C_{3-6} cycloalkyl, C_{2-5} alkenyl, C_{2-5} alkynyl, heterocycloalkyl, aryl and aryl C_{1-4} alkyl; or R_1 and R_1' together form a 3 to 7 membered optionally substituted heterocyclic ring; wherein
- 10 any substituents are selected from the group consisting of CN, aryl, CO_2R , CO_2NHR , OH, OR, NH_2 , halo, CF_3 , OCF_3 and NO_2 ; wherein R represents C_{1-4} alkyl, or C_{3-6} cycloalkyl;
- X_1 is selected from the group consisting of CN, NO_2 , Cl, F, Br, I, H, R', OR', CF_3 , OCF_3 and OSO_2R' , wherein R' represents C_{1-4} alkyl, or C_{3-6} cycloalkyl;
- 15 X_2 , X_3 and X_4 are, independently, selected from the group consisting of CN, NO_2 , Cl, F, Br, I, H, R'', OR'', CF_3 , OCF_3 and OSO_2R'' , provided that either X_1 or X_3 is H, wherein R'' is C_{1-4} alkyl or haloalkyl; or X_1 and X_2 together form an aryl or heteroaryl ring, substituted or unsubstituted; wherein the heteroatom is selected from N, S and O; and any substituents are selected from the group consisting of
- 20 halo, C_{1-4} alkyl, OCF_3 , CF_3 , OMe, CN, OSO_2R' and NO_2 ; or X_3 and X_4 independently represent $C(O)R_1$; and
- R_2 is selected from the group consisting of hydrogen, C_{1-4} alkyl, C_{3-6} cycloalkyl, C_{2-5} alkenyl, C_{2-5} alkynyl, heterocycloalkyl aryl and aryl- C_{1-4} alkyl;
- X_1'' is selected from the group consisting of CN, NO_2 , Cl, F, Br, I, H, R, OR, CF_3 , OCF_3 and OSO_2R , wherein R represents C_{1-4} alkyl, or C_{3-6} cycloalkyl;
- 25 X_2'' , X_3'' and X_4'' are, independently, selected from the group consisting of CN, NO_2 , Cl, F, Br, I, H, R', OR', CF_3 , OCF_3 and OSO_2R' , provided that either X_1'' or X_3'' is H, wherein R' is C_{1-4} alkyl or haloalkyl; or X_1'' and X_2'' together form an aryl or heteroaryl ring, substituted or unsubstituted; wherein the heteroatom is

selected from N, S and O and any substituents are selected from the group consisting of halo, C₁₋₄ alkyl, OCF₃, CF₃, OMe, CN, OSO₂-C₁₋₄ alkyl, OSO₂-C₃₋₆ cycloalkyl and NO₂;

or X₃" and X₄" independently represent C(O)R₁; and

- 5 R₁" and R₂" are, independently, selected from the group consisting of hydrogen, C₁₋₄ alkyl, C₃₋₆ cycloalkyl, C₂₋₅ alkenyl, C₂₋₅ alkynyl, heterocycloalkyl and aryl; or R₁" and R₂" together form a 3 to 7 membered optionally substituted heterocyclic ring; wherein any substituents are selected from the group consisting of CN, aryl, CO₂R", CO₂NHR", OH, OR", NH₂, halo, CF₃, OCF₃ and NO₂;

- 10 wherein R" represents C₁₋₄ alkyl, or C₃₋₆ cycloalkyl;

X₁" is selected from the group consisting of CN, NO₂, Cl, F, Br, I, H, R, OR, CF₃, OCF₃ and OSO₂R, wherein R represents C₁₋₄ alkyl, or C₃₋₆ cycloalkyl;

X₂", X₃", and X₄" are, independently, selected from the group consisting of CN, NO₂, Cl, F, Br, I, H, R', OR', CF₃, OCF₃ and OSO₂R', provided that either X₁" or

- 15 X₃" is H, wherein R' is C₁₋₄ alkyl or haloalkyl;

or X₁" and X₂" together form an aryl or heteroaryl ring, substituted or

unsubstituted; wherein the heteroatom is selected from N, S and O and the

substituents are selected from the group consisting of halo, C₁₋₄ alkyl, OCF₃, CF₃, OMe, CN, OSO₂-C₁₋₄ alkyl, OSO₂-C₃₋₆ cycloalkyl and NO₂;

- 20 or X₃" and X₄" independently represent C(O)R₁;

R₁" and R₂" are, independently, selected from the group consisting of hydrogen,

C₁₋₄ alkyl, C₃₋₆ cycloalkyl, C₂₋₅ alkenyl, C₂₋₅ alkynyl, heterocycloalkyl and aryl;

or R₁" and R₂" together form a 3 to 7 membered optionally substituted

heterocyclic ring; wherein the substituents are selected from the group consisting of

- 25 CN, aryl, CO₂R", CO₂NHR", OH, OR", NH₂, halo, CF₃, OCF₃ and NO₂; wherein R" represents C₁₋₄ alkyl, or C₃₋₆ cycloalkyl;

D is selected from the group consisting of H, CN, NO₂, Cl, F, Br, I, R, OR, SR,

CF₃, OCF₃ and OSO₂R, wherein R represents C₁₋₄ alkyl, C₃₋₆ cycloalkyl, or C₁₋

10 aryl or heteroaryl wherein the heteroatom is selected from N, S and O and

- 30 substituents are selected from the group consisting of halo, C₁₋₄ alkyl, OCF₃, CF₃, OMe, CN, OSO₂-C₁₋₄ alkyl, OSO₂-

C₃₋₆ cycloalkyl and NO₂;

n is the integer 1 or 2;

each E is independently C or N, provided that no more than two E moieties are N;
further provided that when n is 2, each E is C;

a and b are optionally present bonds;

R_1^{IV} is selected from the group consisting of $(CH_2)_nCO_2R'$, $(CH_2)_nCO_2H$,

- 5 $(CH_2)_nCONR'_2$, $(CH_2)_nCH_2OR'$, OR' , SR' , CN , NO_2 , Cl , F , Br , I , H , CF_3 , OCF_3 , OSO_2R' , R' and H ; wherein R' represents C_{1-4} alkyl, or C_{3-6} cycloalkyl;
or R_1^{IV} is O, forming a ketone such that $Y R_1^{IV}$ represents $-C=O$;

R_2^{IV} is selected from the group consisting of hydrogen, CN , NO_2 , Cl , F , Br , I , H , R'' , OR'' , CF_3 , OCF_3 , and OSO_2R'' ; wherein R'' represents C_{1-4} alkyl, or C_{3-6}

- 10 cycloalkyl.

Y is selected from the group consisting of C , CH , O , N and S ; provided that when Y is S , R_1^{IV} is O or not present; further provided that when Y is O , R_1^{IV} is not present;

X' is selected from the group consisting of CH_2 , NH , O and S .

- 15 R_9 is selected from the group consisting of O -alkyl, O - CH_2 -aryl, and O -aryl;

X_1^{III} is selected from the group consisting of CN , NO_2 , Cl , F , Br , I , H , R , OR , CF_3 , OCF_3 and OSO_2R , wherein R represents C_{1-4} alkyl, or C_{3-6} cycloalkyl;

X_2^{III} , X_3^{III} , and X_4^{III} are, independently, selected from the group consisting of CN , NO_2 , Cl , F , Br , I , H , R' , OR' , CF_3 , OCF_3 and OSO_2R' , provided that either X^{III}_1 or


- 20 X^{III}_3 is H , wherein R' is C_{1-4} alkyl or haloalkyl;

or X_1^{III} and X_2^{III} together form an aryl or heteroaryl ring, substituted or unsubstituted; wherein the heteroatom is selected from N , S and O and the substituents are selected from the group consisting of halo, C_{1-4} alkyl, OCF_3 , CF_3 , OMe , CN , OSO_2 - C_{1-4} alkyl, OSO_2 - C_{3-6} cycloalkyl and NO_2 ;

- 25 or X_2^{III} and X_4^{III} independently represent $C(O)R_1$;

and pharmaceutically acceptable salts and complexes thereof.

2. A compound according to claim 1 having a structure according to Formula (II) hereinbelow::



Formula (II)

30

wherein:

R_5 is heteroaryl or fused heteroaryl; wherein the hetero-ring contains N, O or S, and is aromatic, dihydro or tetrahydro, unsubstituted or substituted with any substituents being selected from the group consisting of OH, OCH_3 , $CH(CH_3)_2$, halogen, C_{1-4} alkyl, C_{1-4} alkoxy, C_{3-6} cycloalkyl, OSO_2R^{IV} , CN, NO_2 , OCF_3 , CF_3 , CH_2CF_3 , $(CH_2)_n CO_2H$, $(CH_2)_n CO_2R^{IV}$, and $O-(CH_2)_n CO_2R^{IV}$; and
 5 A and B are, independently, selected from the group consisting of a bond, CH_2 , NH, O, S and $C=O$, provided that either A or B is selected from CH_2 and NH; or A and B together form a bond; or the A-B moiety is represented by $CH=CH$ or $C\equiv C$.

3. A compound according to claim 2 wherein:

10 R_5 is heteroaryl or fused heteroaryl, wherein the hetero-ring contains N, O or S and is aromatic, dihydro or tetrahydro, unsubstituted or substituted with any substituents being selected from the group consisting of OCH_3 , halogen, C_{1-4} alkyl, , CN, NO_2 , OCF_3 , CF_3 , and CH_2CF_3 ;

R_6 is H; and

15 A and B are, independently, selected from the group consisting of a bond, CH_2 , NH, O, S and $C=O$, provided that either A or B is selected from CH_2 and NH, or A and B together form a bond.

4. A compound according to claim 3 wherein:

R_5 is heteroaryl or fused heteroaryl, wherein the hetero-ring contains N, O or S and
 20 is aromatic, dihydro or tetrahydro, unsubstituted or substituted with any substituents being selected from the group consisting of OCH_3 , halogen, C_{1-4} alkyl, , CN, NO_2 , OCF_3 , CF_3 , and CH_2CF_3 ;

R_6 is H; and

A and B are, independently, selected from the group consisting of a bond, CH_2 , O,
 25 or A and B together form a bond.

5. A compound according to claim 1 selected from the group consisting of:

(R)-N-[2-Hydroxy-3-(3-chloro-2-cyanophenoxy)propyl]-1,1-dimethyl-2-(2,3-dihydrobenzo[b]furan-5yl)ethylamine;

(R)-N-[2-Hydroxy-3-(3-chloro-2-cyanophenoxy)propyl]-1,1-dimethyl-2-(quinolin-
 30 3-yl)ethylamine;

(R)-N-[2-Hydroxy-3-(3-chloro-2-cyanophenoxy)propyl]-1,1-dimethyl-2-(quinolin-2-yl)ethylamine;

- (R)-N-[2-Hydroxy-3-(3-chloro-2-cyanophenoxy)propyl]-1,1-dimethyl-2-(isoquinolin-3-yl)ethylamine;
- (R)-N-[2-Hydroxy-3-(3-chloro-2-cyanophenoxy)propyl]-1,1-dimethyl-4-(2-pyridyl)butylamine;
- 5 (R)-N-[2-Hydroxy-3-(3-chloro-2-cyano-4-morpholinosulfonamidophenoxy)propyl]-1,1-dimethyl-4-(2-pyridyl)butylamine;
- (R)-N-[2-Hydroxy-3-(3-chloro-2-cyanophenoxy)propyl]-1,1-dimethyl-4-(3-pyridyl)butylamine;
- (R)-N-[2-Hydroxy-3-(3-chloro-2-cyano-4-morpholinosulfonamidophenoxy)propyl]-1,1-dimethyl-4-(3-pyridyl)butylamine;
- 10 (R)-N-[2-Hydroxy-3-(3-chloro-2-cyanophenoxy)propyl]-1,1-dimethyl-4-(4-carbethoxyphenyl)butylamine;
- (R)-N-[2-Hydroxy-3-(3-chloro-2-cyanophenoxy)propyl]-1,1-dimethyl-2-(4-ethylpyrid-2-yl)ethylamine;
- 15 (R)-N-[2-Hydroxy-3-(3-chloro-2-cyanophenoxy)propyl]-1,1-dimethyl-2-benzamidoethylamine;
- (R)-N-[2-Hydroxy-3-(3-chloro-2-cyanophenoxy)propyl]-4-phenylbutylamine;
- (R)-N-[2-Hydroxy-3-(3-chloro-2-cyanophenoxy)propyl]-1,1-dimethyl-4-phenylbut-2-ynylamine;
- 20 and pharmaceutically acceptable salts and complexes thereof.
6. A compound according to claim 5 selected from the group consisting of:
- (R)-N-[2-Hydroxy-3-(3-chloro-2-cyanophenoxy)propyl]-1,1-dimethyl-2-(2,3-dihydrobenzo[b]furan-5-yl)ethylamine;
- (R)-N-[2-Hydroxy-3-(3-chloro-2-cyanophenoxy)propyl]-1,1-dimethyl-2-(quinolin-
- 25 3-yl)ethylamine;
- (R)-N-[2-Hydroxy-3-(3-chloro-2-cyanophenoxy)propyl]-1,1-dimethyl-2-(quinolin-2-yl)ethylamine;
- (R)-N-[2-Hydroxy-3-(3-chloro-2-cyanophenoxy)propyl]-1,1-dimethyl-2-(isoquinolin-3-yl)ethylamine;
- 30 (R)-N-[2-Hydroxy-3-(3-chloro-2-cyanophenoxy)propyl]-1,1-dimethyl-4-(2-pyridyl)butylamine;
- (R)-N-[2-Hydroxy-3-(3-chloro-2-cyano-4-morpholinosulfonamidophenoxy)propyl]-1,1-dimethyl-4-(2-pyridyl)butylamine;

(R)-N-[2-Hydroxy-3-(3-chloro-2-cyanophenoxy)propyl]-1,1-dimethyl-4-(3-pyridyl)butylamine;

(R)-N-[2-Hydroxy-3-(3-chloro-2-cyano-4-morpholinosulfonamidophenoxy)propyl]-1,1-dimethyl-4-(3-pyridyl)butylamine;

5 (R)-N-[2-Hydroxy-3-(3-chloro-2-cyanophenoxy)propyl]-1,1-dimethyl-2-(4-ethylpyrid-2-yl)ethylamine;

(R)-N-[2-Hydroxy-3-(3-chloro-2-cyanophenoxy)propyl]-1,1-dimethyl-2-benzamidoethylamine; and

(R)-N-[2-Hydroxy-3-(3-chloro-2-cyanophenoxy)propyl]-4-phenylbutylamine;

10 and pharmaceutically acceptable salts and complexes thereof.

7. A compound according to claim 6 selected from the group consisting of:

(R)-N-[2-Hydroxy-3-(3-chloro-2-cyanophenoxy)propyl]-1,1-dimethyl-2-(2,3-dihydrobenzo[b]furan-5yl)ethylamine;

15 (R)-N-[2-Hydroxy-3-(3-chloro-2-cyanophenoxy)propyl]-1,1-dimethyl-2-(quinolin-3-yl)ethylamine;

(R)-N-[2-Hydroxy-3-(3-chloro-2-cyanophenoxy)propyl]-1,1-dimethyl-2-(quinolin-2-yl)ethylamine;

(R)-N-[2-Hydroxy-3-(3-chloro-2-cyanophenoxy)propyl]-1,1-dimethyl-2-(isoquinolin-3-yl)ethylamine;

20 (R)-N-[2-Hydroxy-3-(3-chloro-2-cyanophenoxy)propyl]-1,1-dimethyl-4-(2-pyridyl)butylamine;

(R)-N-[2-Hydroxy-3-(3-chloro-2-cyano-4-morpholinosulfonamidophenoxy)propyl]-1,1-dimethyl-4-(2-pyridyl)butylamine;

25 (R)-N-[2-Hydroxy-3-(3-chloro-2-cyanophenoxy)propyl]-1,1-dimethyl-4-(3-pyridyl)butylamine;

(R)-N-[2-Hydroxy-3-(3-chloro-2-cyano-4-morpholinosulfonamidophenoxy)propyl]-1,1-dimethyl-4-(3-pyridyl)butylamine;

and


30 (R)-N-[2-Hydroxy-3-(3-chloro-2-cyanophenoxy)propyl]-1,1-dimethyl-2-(4-ethylpyrid-2-yl)ethylamine;

and pharmaceutically acceptable salts and complexes thereof.

8. A pharmaceutical composition for use in treating a disease or disorder characterized by an abnormal bone or mineral homeostasis which comprises a compound according to claim 1 and a pharmaceutically acceptable carrier.
9. A method of antagonizing a calcium receptor which comprises
5 administering to a subject in need thereof an effective amount of a compound according to claim 1.
10. A method of treating a disease or disorder characterized by an abnormal bone or mineral homeostasis which comprises administering to a subject in need of treatment an effective amount of a compound according to claim 1.
- 10 11. A method according to claim 10 wherein the bone or mineral disease or disorder is selected from the group consisting of osteosarcoma, periodontal disease, fracture healing, osteoarthritis, rheumatoid arthritis, Paget's disease, humoral hypercalcemia, malignancy and osteoporosis.
12. A method according to claim 11 wherein the bone or mineral disease or
15 disorder is osteoporosis.
13. A method of increasing serum parathyroid levels which comprises administering to a subject in need of treatment an effective amount of a compound according to claim 1.
14. Use of a compound according to claim 1 in the manufacture of a
20 medicament for use in treating a disease or disorder characterized by an abnormal bone or mineral homeostasis.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US99/07760

A. CLASSIFICATION OF SUBJECT MATTER IPC(6) : Please See Extra Sheet. US CL : Please See Extra Sheet. According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) U.S. : Please See Extra Sheet. Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) CAS ONLINE. APS		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X, P	Chem. abstr., No. 302376, Vol. 129, BARMORE et al, 'Preparation of arylalkylamine as calcilytic compounds,' abstract, WO 98/45255 15 October 1198, see abstract.	1-14
Y	US 5,693,310 A (GRIES et al) 02 December 1997, col. 2, lines 15-55.	1-14
<input type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.		
* *A* *B* *L* *O* *P*	Special categories of cited documents: document defining the general state of the art which is not considered to be of particular relevance earlier document published on or after the international filing date document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) document referring to an oral disclosure, use, exhibition or other means document published prior to the international filing date but later than the priority date claimed	*T* *X* *Y* *Z* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art document member of the same patent family
Date of the actual completion of the international search 03 JULY 1999		Date of mailing of the international search report 19 AUG 1999
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 Facsimile No. (703) 305-3230		Authorized officer RAYMOND COVINGTON  Telephone No. (703) 308-1235

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US99/07760

A. CLASSIFICATION OF SUBJECT MATTER:

IPC (6):

A61K 31/535; A01N 43/02, 43/40, 43/42; C07D 211/70, 211/72, 217/12, 217/16, 217/18, 217/38, 217/60, 307/02

A. CLASSIFICATION OF SUBJECT MATTER:

US CL :

514/237.5, 237.8, 238.2, 307, 309, 310, 311, 312, 131, 345, 469, 470; 544/162, 168, 169; 546/139, 143, 146, 149, 153, 159, 160, 168, 174, 176, 177, 304, 329, 334; 549/467, 480, 491, 493, 494

B. FIELDS SEARCHED

Minimum documentation searched

Classification System: U.S.

514/237.5, 237.8, 238.2, 307, 309, 310, 311, 312, 131, 345, 469, 470; 544/162, 168, 169; 546/139, 143, 146, 149, 153, 159, 160, 168, 174, 176, 177, 304, 329, 334; 549/467, 480, 491, 493, 494